arXiv:1003.5882v1 [math.QA] 30 Mar 2010

On the lifting of Nichols algebras*

Michael Helbig†

March 31, 2010

Abstract

Nichols algebras are a fundamental building block of pointed Hopf algebras. Part of the classification program of finite-dimensional pointed Hopf algebras with the lifting method of Andruskiewitsch and Schneider is the determination of the liftings, i.e., all possible deformations of a given Nichols algebra. Based on recent work of Heckenberger about Nichols algebras of diagonal type we compute explicitly the liftings of

- all Nichols algebras with Cartan matrix of type A_2,
- some Nichols algebras with Cartan matrix of type B_2, and
- some Nichols algebras of two Weyl equivalence classes of non-standard type

giving new classes of finite-dimensional pointed Hopf algebras.

Key Words: Hopf algebra, Nichols algebra, quantum group, lifting

Introduction

At the moment the most promissing general method for the classification of finite-dimensional pointed Hopf algebras is the lifting method developed by Andruskiewitsch and Schneider [3]: Given a finite-dimensional pointed Hopf algebra A with coradical $A_0 = \Bbbk[\Gamma]$ and abelian group of group-like elements $\Gamma = G(A)$. Then we can decompose its associated graded Hopf algebra into a smash product $\mathrm{gr}(A) \cong B\#\Bbbk[\Gamma]$ where B is a braided Hopf algebra. The subalgebra of B generated by its primitive elements $V := P(B)$ is a Nichols algebra $\mathfrak{B}(V)$. Now the classification is carried out in three steps:

(1) Show that $B = \mathfrak{B}(V)$.

(2) Determine the structure of $\mathfrak{B}(V)$.

(3) *Lifting*: Determine the liftings of $\mathfrak{B}(V)$, i.e., all Hopf algebras A such that $\mathrm{gr}(A) \cong \mathfrak{B}(V)\#\Bbbk[\Gamma]$.

Many classification results in special situations were obtained in this way [3, 5, 4, 12, 13]. The most impressive result obtained by this method by Andruskiewitsch und Schneider [7] is the classification of all finite-dimensional pointed Hopf algebras where the prime divisors of the order of the abelian group Γ are > 7. In this case the diagonal braiding of V is of

---

[*] This work is part of the author's PhD thesis written under the supervision of Professor H.-J. Schneider.

[†] eMail: `michael@helbig123.de`

Cartan type and the Hopf algebras are generalized versions of small quantum groups. The classification when the braiding is not of Cartan type or the divisors of the order of Γ are ≤ 7 is still an open problem. Also the case where Γ is not abelian is widely open and of different nature, e.g., the defining relations have another form [20, 2].

Concerning (2), Heckenberger recently showed that Nichols algebras of diagonal type have a close connection to semi-simple Lie algebras, namely he introduces a Weyl groupoid [16], Weyl equivalence [15] and an arithmetic root system [18, 14] for Nichols algebras. With the help of these concepts he classifies the diagonal braidings of V such that the Nichols algebra $\mathfrak{B}(V)$ has a finite set of PBW generators [19]. Moreover, he determines the structure of all rank two Nichols algebras in terms of generators and relations [17].

This is the starting point of our work which addresses to step (3) of the program, namely the *lifting* in the cases not treated in [7]. As said before, we lift Nichols algebras of diagonal type with Cartan matrix of type A_2 in Theorem 5.9, with Cartan matrix of type B_2 in Theorem 5.13, and of non-standard type in Theorem 5.17: When lifting arbitrary diagonal Nichols algebras, new phenomena occur: In the setting of [7] there are only three types of defining relations, namely the Serre relations, the linking relations and the root vector relations. The algebraic structure in the general setting is more complicated: Firstly, the Serre relations do not play the outstanding role. Other relations are needed and sometimes the Serre relations are redundant; we give a complete answer for the Serre relations in Lemma 5.6. Secondly, in general the lifted relations from the Nichols algebra do not remain in the group algebra.

The paper is organized as follows: In Section 1 we recall the basic notions of Nichols algebras of diagonal type, taking into account the recent developement by Heckenberger. Then in Section 2 the general calculus for q-commutators in an arbitrary algebra of [21, 23] is presented. Section 3 contains the theory of Lyndon words, super letters and super words. Crucial for the lifting is the knowledge of a "good" presentation of the Nichols algebra and its liftings, in terms of generators and (non-redundant) relations: Section 4 repeats the result of [22]; further some new formulas for coproducts are developed. We then formulate the main results about the liftings in Section 5. We explain our method in Section 5.1.

Throughout the paper $\Bbbk$ will be a field, all vector spaces will be over $\Bbbk$, and all tensor products are taken over $\Bbbk$.

1 Nichols algebras

We want to define Nichols algebras in the category ${}^{\Gamma}_{\Gamma}\mathcal{YD}$ of Yetter-Drinfel'd modules over an abelian group Γ, not necessarily finite. Our main reference is the survey article [6, Sect. 1,2].

1.1 Yetter-Drinfel'd modules of diagonal type

The category ${}^{\Gamma}_{\Gamma}\mathcal{YD}$ of (left-left) Yetter-Drinfel'd modules over the Hopf algebra $\Bbbk[\Gamma]$ is the category of left $\Bbbk[\Gamma]$-modules which are Γ-graded vector spaces $V = \bigoplus_{g\in\Gamma} V_g$ such that each V_g is stable under the action of Γ, i.e., $h \cdot v \in V_g \quad$ for all $\quad h \in \Gamma, v \in V_g$. The Γ-grading is equivalent to a left $\Bbbk[\Gamma]$-comodule structure $\delta : V \to \Bbbk[\Gamma] \otimes V$: One can define δ or the other way round V_g by the equivalence $\delta(v) = g \otimes v \iff v \in V_g$ for all $g \in \Gamma$. The morphisms of ${}^{\Gamma}_{\Gamma}\mathcal{YD}$ are the Γ-linear maps $f : V \to W$ with $f(V_g) \subset W_g$ for all $g \in \Gamma$.

We consider the following monoidal structure on ${}_{\Gamma}^{\Gamma}\mathcal{YD}$: If $V, W \in {}_{\Gamma}^{\Gamma}\mathcal{YD}$, then also $V \otimes W \in {}_{\Gamma}^{\Gamma}\mathcal{YD}$ by

$$g \cdot (v \otimes w) := (g \cdot v) \otimes (g \cdot w) \quad \text{and} \quad (V \otimes W)_g := \bigoplus_{hk=g} V_h \otimes W_k$$

for $v \in V, w \in W$ and $g \in \Gamma$. The *braiding* in ${}_{\Gamma}^{\Gamma}\mathcal{YD}$ is the isomorphism

$$c = c_{V,W} : V \otimes W \to W \otimes V, \quad c(v \otimes w) := (g \cdot w) \otimes v$$

for all $v \in V_g$, $g \in \Gamma$, $w \in W$. Thus every $V \in {}_{\Gamma}^{\Gamma}\mathcal{YD}$ is a braided vector space $(V, c_{V,V})$.

We have the following important example: For a group Γ we denote by $\widehat{\Gamma}$ the character group of all group homomorphisms from Γ to the multiplicative group $\Bbbk^{\times}$.

Definition 1.1. Let $V \in {}_{\Gamma}^{\Gamma}\mathcal{YD}$. If there is a basis x_i, $i \in I$, of V and $g_i \in \Gamma$, $\chi_i \in \widehat{\Gamma}$ for all $i \in I$ such that

$$g \cdot x_i = \chi_i(g) x_i \quad \text{and} \quad x_i \in V_{g_i},$$

then we say V is of *diagonal type.*

Note that if $\Bbbk$ is algebraically closed of characteristic 0 and Γ is finite, then any finite-dimensional $V \in {}_{\Gamma}^{\Gamma}\mathcal{YD}$ is of diagonal type.

For the braiding we have $c(x_i \otimes x_j) = \chi_j(g_i) x_j \otimes x_i$ for $1 \leq i, j \leq \theta$. Hence the braiding is determined by the so-called *braiding matrix* of V

$$(q_{ij})_{1 \leq i,j \leq \theta} := (\chi_j(g_i))_{1 \leq i,j \leq \theta}.$$

1.2 Nichols algebras of diagonal type

Let $V \in {}_{\Gamma}^{\Gamma}\mathcal{YD}$. B is called a *Nichols algebra* of V, if

- $B = \oplus_{n \geq 0} B(n)$ is a graded braided Hopf algebra in ${}_{\Gamma}^{\Gamma}\mathcal{YD}$,
- $B(0) \cong \Bbbk$,
- $P(B) = B(1) \cong V$,
- B is generated as an algebra by $B(1)$.

Any two Nichols algebras of V are isomorphic, thus we write $\mathfrak{B}(V)$ for "the" Nichols algebra of V. One can construct the Nichols algebra in the following way: Let I denote the sum of all ideals of $T(V)$ that are generated by homogeneous elements of degree ≥ 2 and that are also coideals. Then $\mathfrak{B}(V) \cong T(V)/I$. We say the Nichols algebra $\mathfrak{B}(V)$ is of *diagonal type*, if V is of diagonal type.

1.3 Cartan matrices

A matrix $(a_{ij})_{1 \leq i,j \leq \theta} \in \mathbb{Z}^{\theta \times \theta}$ is called a *generalized Cartan matrix* if for all $1 \leq i, j \leq \theta$

- $a_{ii} = 2$,
- $a_{ij} \leq 0$ if $i \neq j$,
- $a_{ij} = 0 \Rightarrow a_{ji} = 0$.

Let $\mathfrak{B}(V)$ be a Nichols algebra of diagonal type. Recall that V resp. $\mathfrak{B}(V)$ with braiding matrix (q_{ij}) is called of *Cartan type*, if there is a generalized Cartan matrix (a_{ij}) such that

$$q_{ij}q_{ji} = q_{ii}^{a_{ij}}.$$

Not every Nichols algebra is of Cartan type (see Example 1.3 and Sections 5.3, 5.4, 5.5), but still we have the following [16, Sect. 3]:

If $\mathfrak{B}(V)$ is finite-dimensional, then the matrix (a_{ij}) defined for all $1 \leq i \neq j \leq \theta$ by

$$a_{ii} := 2 \quad \text{and} \quad a_{ij} := -\min\{r \in \mathbb{N} \mid q_{ij}q_{ji}q_{ii}^r = 1 \text{ or } (r+1)_{q_{ii}} = 0\}$$

is a generalized Cartan matrix fulfilling

$$q_{ij}q_{ji} = q_{ii}^{a_{ij}} \quad \text{or} \quad \text{ord} q_{ii} = 1 - a_{ij}. \tag{1.1}$$

We call (a_{ij}) the *Cartan matrix associated to* $\mathfrak{B}(V)$.

1.4 Weyl equivalence

Heckenberger introduced in [15, 16, Sect. 2] the notion of the *Weyl groupoid* and *Weyl equivalence* of Nichols algebras of diagonal type. With the help of these concepts Heckenberger classified in a series of articles [18, 14, 19] all braiding matrices (q_{ij}) of diagonal Nichols algebras with a finite set of PBW generators. We are concerned with the list of rank 2 Nichols algebras given in the table [15, 18, Figure 1].

We want to recall the following: For diagonal $\mathfrak{B}(V)$ with braiding matrix (q_{ij}) we associate a *generalized Dynkin diagram*: this is a graph with θ vertices, where the i-th vertex is labeled with q_{ii} for all $1 \leq i \leq \theta$; further, if $q_{ij}q_{ji} \neq 1$, then there is an edge between the i-th and j-th vertex labeled with $q_{ij}q_{ji}$: Thus, if $q_{ij}q_{ji} = 1$ resp. $q_{ij}q_{ji} \neq 1$, then we have

resp.



So two Nichols algebras of the same rank θ with braiding matrix (q_{ij}) resp. (q'_{ij}) have the same generalized Dynkin diagram if and only if they are *twist equivalent* [6, Def. 3.8], i.e., for all $1 \leq i, j \leq \theta$

$$q_{ii} = q'_{ii} \quad \text{and} \quad q_{ij}q_{ji} = q'_{ij}q'_{ji}.$$

Definition 1.2. Let $1 \leq k \leq \theta$ be fixed and $\mathfrak{B}(V)$ finite-dimensional with braiding matrix (q_{ij}) and Cartan matrix (a_{ij}). We call $(q_{ij}^{(k)})$ defined by

$$q_{ij}^{(k)} := q_{ij}q_{ik}^{-a_{kj}}q_{kj}^{-a_{ki}}q_{kk}^{a_{ki}a_{kj}}$$

the *at the vertex k reflected braiding matrix.*

Two Nichols algebras with braiding matrix (q_{ij}) resp. (q'_{ij}) are called *Weyl equivalent*, if there are $m \geq 1$, $1 \leq k_1, \ldots, k_m \leq \theta$ such that the generalized Dynkin diagrams w.r.t. the matrices $\big((\ldots(q_{ij}^{(k_1)})^{(k_2)}\ldots)^{(k_m)}\big)$ and (q'_{ij}) coincide, i.e., one gets the Dynkin diagram of (q'_{ij}) by successive reflections of (q_{ij}).

Example 1.3. The braiding matrix $(q_{ij}) := \begin{pmatrix} q & 1 \\ q^{-1} & -1 \end{pmatrix}$ with $q \neq 1$ has the generalized Dynkin diagram $\overset{q}{\circ}\overset{q^{-1}}{\text{———}}\overset{-1}{\circ}$ and associated Cartan matrix $(a_{ij}) = \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix}$ of type A_2, since $q_{12}q_{21} = q_{11}^{-1}$ and $\operatorname{ord} q_{22} = 1 - (-1)$. Then the at the vertex 2 reflected braiding matrix is $(q'_{ij}) := (q_{ij}^{(2)}) = \begin{pmatrix} -1 & -1 \\ -q & -1 \end{pmatrix}$. Its Dynkin diagram is $\overset{-1}{\circ}\overset{q}{\text{———}}\overset{-1}{\circ}$ and the associated Cartan matrix is also of type A_2. The two braiding matrices (q_{ij}) and (q'_{ij}) are by definition Weyl equivalent; they are twist equivalent if and only if $q = -1$. See also [15, 18, Figure 1]: row 3 if $q \neq \pm 1$ and row 2 if $q = -1$.

Remark 1.4.

1. Both twist equivalence and Weyl equivalence are equivalence relations, and twist equivalent Nichols algebras are Weyl equivalent.

2. Weyl equivalent Nichols algebras have the same dimension and Gel'fand-Kirillov dimension [15, Prop. 1], but can have different associated Cartan matrices. If the whole Weyl equivalence class has the same Cartan matrix, then the Nichols algebras of this class are called of *standard* type [1, 8].

Examples 1.5. Let $\mathfrak{B}(V)$ be of rank 2. Then two Nichols algebras are Weyl equivalent if and only if their generalized Dynkin diagrams appear in the same row of [15, 18, Figure 1] and can be presented with the same set of fixed parameters [15].

1. $\mathfrak{B}(V)$ is of standard type, if and only if it appears in the rows 1–7, 11 or 12 of [15, 18, Figure 1]. The Cartan matrices are

   - $\begin{pmatrix} 2 & 0 \\ 0 & 2 \end{pmatrix}$ of type $A_1 \times A_1$ of row 1,
   - $\begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix}$ of type A_2 of rows 2 and 3,
   - $\begin{pmatrix} 2 & -2 \\ -1 & 2 \end{pmatrix}$ of type B_2 of rows 4–7, and
   - $\begin{pmatrix} 2 & -3 \\ -1 & 2 \end{pmatrix}$ of type G_2 of rows 11 and 12.

   All Nichols algebras of type $A_1 \times A_1$, A_2 and some of B_2 are lifted in Sections 5.2, 5.3, 5.4.

2. In the non-standard Weyl equivalence class of row 8 of [15, 18, Figure 1] the Cartan matrices

   - $\begin{pmatrix} 2 & -2 \\ -2 & 2 \end{pmatrix}$ of $\overset{-\zeta^{-2}}{\circ}\overset{-\zeta^{3}}{\text{———}}\overset{-\zeta^{2}}{\circ}$,
   - $\begin{pmatrix} 2 & -2 \\ -1 & 2 \end{pmatrix}$ of type B_2 of $\overset{-\zeta^{-2}}{\circ}\overset{\zeta^{-1}}{\text{———}}\overset{-1}{\circ}$, $\overset{-\zeta^{2}}{\circ}\overset{-\zeta}{\text{———}}\overset{-1}{\circ}$, and
   - $\begin{pmatrix} 2 & -3 \\ -1 & 2 \end{pmatrix}$ of type G_2 of $\overset{-\zeta^{3}}{\circ}\overset{\zeta}{\text{———}}\overset{-1}{\circ}$, $\overset{-\zeta^{3}}{\circ}\overset{-\zeta^{-1}}{\text{———}}\overset{-1}{\circ}$

   appear. These Nichols algebras are lifted in Section 5.5. The same Cartan matrices appear in row 9, where we lift the Nichols algebras corresponding to the last two Dynkin diagrams.

2 q-commutator calculus

In this section let A denote an arbitrary algebra over a field $\Bbbk$ of characteristic $\operatorname{char} \Bbbk = p \geq 0$. The main result of this chapter is Proposition 2.1, which states important q-commutator formulas in an arbitrary algebra.

2.1 q-calculus

For every $q \in \Bbbk$ we define for $n \in \mathbb{N}$ and $0 \leq i \leq n$ the *q-numbers* $(n)_q := 1 + q + q^2 + \ldots + q^{n-1}$, the *$q$-factorials* $(n)_q! := (1)_q(2)_q \ldots (n)_q$, and the *$q$-binomial coefficients* $\binom{n}{i}_q := \frac{(n)_q!}{(n-i)_q!(i)_q!}$. Note that the latter right-handside is well-defined since it is a polynomial over $\mathbb{Z}$ evaluated in q. We denote the *multiplicative order* of any $q \in \Bbbk^\times$ by $\mathrm{ord} q$. If $q \in \Bbbk^\times$ and $n > 1$, then

$$\binom{n}{i}_q = 0 \text{ for all } 1 \leq i \leq n-1 \Longleftrightarrow \begin{cases} \mathrm{ord} q = n, & \text{if } \operatorname{char} \Bbbk = 0 \\ p^k \mathrm{ord} q = n \text{ with } k \geq 0, & \text{if } \operatorname{char} \Bbbk = p > 0, \end{cases} \tag{2.1}$$

see [27, Cor. 2]. Moreover for $1 \leq i \leq n$ there are the *q-Pascal identities*

$$q^i \binom{n}{i}_q + \binom{n}{i-1}_q = \binom{n}{i}_q + q^{n+1-i} \binom{n}{i-1}_q = \binom{n+1}{i}_q, \tag{2.2}$$

and the *q-binomial theorem*: For $x, y \in A$ and $q \in \Bbbk^\times$ with $yx = qxy$ we have

$$(x+y)^n = \sum_{i=0}^{n} \binom{n}{i}_q x^i y^{n-i}. \tag{2.3}$$

Note that for $q = 1$ these are the usual notions.

2.2 q-commutators

Let $\theta \geq 1$, $X = \{x_1, \ldots, x_\theta\}$, $\langle X \rangle$ the free monoid and $A = \Bbbk\langle X \rangle$ the free $\Bbbk$-algebra. For an abelian group Γ let $\widehat{\Gamma}$ be the character group, $g_1, \ldots, g_\theta \in \Gamma$ and $\chi_1, \ldots, \chi_\theta \in \widehat{\Gamma}$. If we define the two monoid maps

$$\deg_\Gamma : \langle X \rangle \to \Gamma,\ \deg_\Gamma(x_i) := g_i \quad \text{and} \quad \deg_{\widehat{\Gamma}} : \langle X \rangle \to \widehat{\Gamma},\ \deg_{\widehat{\Gamma}}(x_i) := \chi_i,$$

for all $1 \leq i \leq \theta$, then $\Bbbk\langle X \rangle$ is Γ- and $\widehat{\Gamma}$-graded. Let $a \in \Bbbk\langle X \rangle$ be Γ-homogeneous and $b \in \Bbbk\langle X \rangle$ be $\widehat{\Gamma}$-homogeneous. We set

$$g_a := \deg_\Gamma(a), \quad \chi_b := \deg_{\widehat{\Gamma}}(b), \quad \text{and} \quad q_{a,b} := \chi_b(g_a).$$

Further we define $\Bbbk$-linearly on $\Bbbk\langle X \rangle$ the *q-commutator*

$$[a, b] := [a, b]_{q_{a,b}}. \tag{2.4}$$

Note that $q_{a,b}$ is a bicharacter on the homogeneous elements and depends only on the values

$$q_{ij} := \chi_j(g_i) \text{ with } 1 \leq i, j \leq \theta.$$

For example $[x_1, x_2] = x_1x_2 - \chi_2(g_1)x_2x_1 = x_1x_2 - q_{12}x_2x_1$. Further if a, b are $\mathbb{Z}^\theta$-homogeneous they are both Γ- and $\widehat{\Gamma}$-homogeneous. In this case we can build iterated q-commutators, like $\big[x_1, [x_1, x_2]\big] = x_1[x_1, x_2] - \chi_1\chi_2(g_1)[x_1, x_2]x_1 = x_1[x_1, x_2] - q_{11}q_{12}[x_1, x_2]x_1$.

Proposition 2.1. [23, Prop. 1.2] *For all homogeneous $a, b, c \in \Bbbk\langle X\rangle$ and $r \geq 1$ we have:*

(1) q-derivation properties: $[a, bc] = [a, b]c + q_{a,b}b[a, c]$, $[ab, c] = a[b, c] + q_{b,c}[a, c]b$.

(2) q-Jacobi identity: $\big[[a, b], c\big] = \big[a, [b, c]\big] - q_{a,b}b[a, c] + q_{b,c}[a, c]b$.

(3) q-Leibniz formulas:

$$[a, b^r] = \sum_{i=0}^{r-1} q_{a,b}^i \binom{r}{i}_{q_{b,b}} b^i \big[\dots \big[[a, \underbrace{b], b\big] \dots, b}_{r-i}\big],$$

$$[a^r, b] = \sum_{i=0}^{r-1} q_{a,b}^i \binom{r}{i}_{q_{a,a}} \underbrace{\big[a, \dots \big[a, [a}_{r-i}, b]\big] \dots\big] a^i.$$

(4) restricted q-Leibniz formulas: *If* $\operatorname{char}\Bbbk = 0$ *and* $\operatorname{ord}q_{b,b} = r$ *resp.* $\operatorname{ord}q_{a,a} = r$*, or* $\operatorname{char}\Bbbk = p > 0$ *and* $p^k\operatorname{ord}q_{b,b}\zeta = r$ *resp.* $p^k\operatorname{ord}q_{a,a} = r$*, then*

$$[a, b^r] = \big[\dots \big[[a, \underbrace{b], b\big] \dots, b}_{r}\big] \quad \textit{resp.} \quad [a^r, b] = \big[\underbrace{a, \dots \big[a, [a}_{r}, b]\big] \dots\big].$$

3 Lyndon words and q-commutators

In this section we recall the theory of Lyndon words [25, 28] as far as we are concerned and then introduce the notion of super letters and super words [24].

3.1 Words and the lexicographical order

Let $\theta \geq 1$, $X = \{x_1, x_2, \dots, x_\theta\}$ be a finite totally ordered set by $x_1 < x_2 < \dots < x_\theta$, and $\langle X\rangle$ the free monoid; we think of X as an alphabet and of $\langle X\rangle$ as the words in that alphabet including the empty word 1. For a word $u = x_{i_1} \dots x_{i_n} \in \langle X\rangle$ we define $\ell(u) := n$ and call it the *length* of u.

The *lexicographical order* $\leq$ on $\langle X\rangle$ is defined for $u, v \in \langle X\rangle$ by $u < v$ if and only if either v begins with u, i.e., $v = uv'$ for some $v' \in \langle X\rangle\backslash\{1\}$, or if there are $w, u', v' \in \langle X\rangle$, $x_i, x_j \in X$ such that $u = wx_iu'$, $v = wx_jv'$ and $i < j$. E.g., $x_1 < x_1x_2 < x_2$.

3.2 Lyndon words and the Shirshov decomposition

A word $u \in \langle X\rangle$ is called a *Lyndon word* if $u \neq 1$ and u is smaller than any of its proper endings, i.e., for all $v, w \in \langle X\rangle\backslash\{1\}$ such that $u = vw$ we have $u < w$. We denote by

$$\mathcal{L} := \{u \in \langle X\rangle \mid u \text{ is a Lyndon word}\}$$

the set of all Lyndon words. For example $X \subset \mathcal{L}$, but $x_i^n \notin \mathcal{L}$ for all $1 \leq i \leq \theta$ and $n \geq 2$. Also x_1x_2, $x_1x_1x_2$, $x_1x_2x_2$, $x_1x_1x_2x_1x_2 \in \mathcal{L}$.

For any $u \in \langle X\rangle\backslash X$ we call the decomposition $u = vw$ with $v, w \in \langle X\rangle\backslash\{1\}$ such that w is the minimal (with respect to the lexicographical order) ending the *Shirshov decomposition* of the word u. We will write in this case

$$\operatorname{Sh}(u) = (v|w).$$

E.g., $\operatorname{Sh}(x_1x_2) = (x_1|x_2)$, $\operatorname{Sh}(x_1x_1x_2x_1x_2) = (x_1x_1x_2|x_1x_2)$, $\operatorname{Sh}(x_1x_1x_2) \neq (x_1x_1|x_2)$. If $u \in \mathcal{L}\backslash X$, this is equivalent to w is the longest proper ending of u such that $w \in \mathcal{L}$.

Definition 3.1. We call a subset $L \subset \mathcal{L}$ *Shirshov closed* if $X \subset L$, and for all $u \in L$ with $\mathrm{Sh}(u) = (v|w)$ also $v, w \in L$.

For example $\mathcal{L}$ is Shirshov closed, and if $X = \{x_1, x_2\}$, then $\{x_1, x_1x_1x_2, x_2\}$ is not Shirshov closed, whereas $\{x_1, x_1x_2, x_1x_1x_2, x_2\}$ is.

3.3 Super letters and super words

Let the free algebra $\Bbbk\langle X\rangle$ be graded as in Section 2.2. For any $u \in \mathcal{L}$ we define recursively on $\ell(u)$ the map

$$[\,.\,] : \mathcal{L} \to \Bbbk\langle X\rangle, \quad u \mapsto [u]. \tag{3.1}$$

If $\ell(u) = 1$, then set $[x_i] := x_i$ for all $1 \leq i \leq \theta$. Else if $\ell(u) > 1$ and $\mathrm{Sh}(u) = (v|w)$ we define $[u] := \big[[v], [w]\big]$. This map is well-defined since inductively all $[u]$ are $\mathbb{Z}^\theta$-homogeneous such that we can build iterated q-commutators; see Section 2.2. The elements $[u] \in \Bbbk\langle X\rangle$ with $u \in \mathcal{L}$ are called *super letters*. E.g. $[x_1x_1x_2x_1x_2] = \big[[x_1x_1x_2], [x_1x_2]\big] = \big[[x_1, [x_1, x_2]], [x_1, x_2]\big]$. If $L \subset \mathcal{L}$ is Shirshov closed then the subset of $\Bbbk\langle X\rangle$

$$[L] := \big\{[u] \,\big|\, u \in L\big\}$$

is a set of iterated q-commutators. Further $[\mathcal{L}] = \big\{[u] \,\big|\, u \in \mathcal{L}\big\}$ is the set of all super letters and the map $[\,.\,] : \mathcal{L} \to [\mathcal{L}]$ is a bijection, which follows from [22, Lem. 2.5]. Hence we can define an order $\leq$ of the super letters $[\mathcal{L}]$ by

$$[u] < [v] :\Leftrightarrow u < v,$$

thus $[\mathcal{L}]$ is a new alphabet containing the original alphabet X; so the name "letter" makes sense. Consequently, products of super letters are called *super words*. We denote

$$[\mathcal{L}]^{(\mathbb{N})} := \big\{[u_1] \ldots [u_n] \,\big|\, n \in \mathbb{N},\, u_i \in \mathcal{L}\big\}$$

the subset of $\Bbbk\langle X\rangle$ of all super words. Any super word has a unique factorization in super letters [22, Prop. 2.6], hence we can define the lexicographical order on $[\mathcal{L}]^{(\mathbb{N})}$, as defined above on regular words. We denote it also by $\leq$.

3.4 A well-founded ordering of super words

The *length* of a super word $U = [u_1][u_2] \ldots [u_n] \in [L]^{(\mathbb{N})}$ is defined as $\ell(U) := \ell(u_1u_2 \ldots u_n)$.

Definition 3.2. For $U, V \in [\mathcal{L}]^{(\mathbb{N})}$ we define $U \prec V$ by

- $\ell(U) < \ell(V)$, or
- $\ell(U) = \ell(V)$ and $U > V$ lexicographically in $[\mathcal{L}]^{(\mathbb{N})}$.

This defines a total ordering of $[\mathcal{L}]^{(\mathbb{N})}$ with minimal element 1. As X is assumed to be finite, there are only finitely many super letters of a given length. Hence every nonempty subset of $[\mathcal{L}]^{(\mathbb{N})}$ has a minimal element, or equivalently, $\preceq$ fulfills the descending chain condition: $\preceq$ is *well-founded*. This makes way for inductive proofs on $\preceq$.

4 A class of pointed Hopf algebras

In this chapter we deal with a special class of pointed Hopf algebras. Let us recall the notions and results of [24, Sect. 3]: A Hopf algebra A is called a *character Hopf algebra* if it is generated as an algebra by elements $a_1, \dots, a_\theta$ and an abelian group $G(A) = \Gamma$ of all group-like elements such that for all $1 \le i \le \theta$ there are $g_i \in \Gamma$ and $\chi_i \in \widehat{\Gamma}$ with

$$\Delta(a_i) = a_i \otimes 1 + g_i \otimes a_i \qquad \text{and} \qquad g a_i = \chi_i(g) a_i g.$$

As mentioned in the introduction this covers a wide class of examples of Hopf algebras.

Theorem 4.1. [22, Thm. 3.4] *If A is a character Hopf algebra, then*

$$A \cong (\Bbbk\langle X\rangle \# \Bbbk[\Gamma])/I,$$

where the smash product $\Bbbk\langle X\rangle \# \Bbbk[\Gamma]$ and the ideal I are constructed in the following way:

4.1 The smash product $\Bbbk\langle X\rangle \# \Bbbk[\Gamma]$

Let $\Bbbk\langle X\rangle$ be Γ- and $\widehat{\Gamma}$-graded as in Section 2.2, and $\Bbbk[\Gamma]$ be endowed with the usual bialgebra structure $\Delta(g) = g \otimes g$ and $\varepsilon(g) = 1$ for all $g \in \Gamma$. Then we define

$$g \cdot x_i := \chi_i(g) x_i, \quad \text{for all } 1 \le i \le \theta.$$

In this case, $\Bbbk\langle X\rangle$ is a $\Bbbk[\Gamma]$-module algebra and we calculate $g x_i = \chi_i(g) x_i g$, $gh = hg = \varepsilon(g) hg$ in $\Bbbk\langle X\rangle \# \Bbbk[\Gamma]$. Thus $x_i \in (\Bbbk\langle X\rangle \# \Bbbk[\Gamma])^{\chi_i}$ and $\Bbbk[\Gamma] \subset (\Bbbk\langle X\rangle \# \Bbbk[\Gamma])^{\varepsilon}$ and in this way $\Bbbk\langle X\rangle \# \Bbbk[\Gamma] = \oplus_{\chi \in \widehat{\Gamma}} (\Bbbk\langle X\rangle \# \Bbbk[\Gamma])^{\chi}$. This $\widehat{\Gamma}$-grading extends the $\widehat{\Gamma}$-grading of $\Bbbk\langle X\rangle$ in Section 2.2 to $\Bbbk\langle X\rangle \# \Bbbk[\Gamma]$. Further $\Bbbk\langle X\rangle \# \Bbbk[\Gamma]$ is a Hopf algebra with structure determined for all $1 \le i \le \theta$ and $g \in \Gamma$ by

$$\Delta(x_i) := x_i \otimes 1 + g_i \otimes x_i \qquad \text{and} \qquad \Delta(g) := g \otimes g.$$

4.2 Ideals associated to Shirshov closed sets

In this subsection we fix a Shirshov closed $L \subset \mathcal{L}$. We want to introduce the following notation for an $a \in \Bbbk\langle X\rangle \# \Bbbk[\Gamma]$ and $W \in [\mathcal{L}]^{(\mathbb{N})}$: We will write $a \prec_L W$ (resp. $a \preceq_L W$), if a is a linear combination of

- $U \in [L]^{(\mathbb{N})}$ with $\ell(U) = \ell(W)$, $U > W$ (resp. $U \ge W$), and
- Vg with $V \in [L]^{(\mathbb{N})}$, $g \in \Gamma$, $\ell(V) < \ell(W)$.

Furthermore, we set for each $u \in L$ either $N_u := \infty$ or $N_u := \operatorname{ord} q_{u,u}$ (resp. $N_u := p^k \operatorname{ord} q_{u,u}$ with $k \ge 0$ if $\operatorname{char} \Bbbk = p > 0$) and we want to distinguish the following two sets of words depending on L:

$$\begin{aligned} C(L) &:= \big\{ w \in \langle X\rangle \backslash L \mid \exists u, v \in L : w = uv,\ u < v, \text{ and } \operatorname{Sh}(w) = (u|v) \big\}, \\ D(L) &:= \big\{ u \in L \mid N_u < \infty \big\}. \end{aligned}$$

Note that $C(L) \subset \mathcal{L}$ and $D(L) \subset L \subset \mathcal{L}$ are sets of Lyndon words. For example, if $L = \{x_1, x_1x_1x_2, x_1x_2, x_2\}$, then $C(L) = \{x_1x_1x_1x_2,\ x_1x_1x_2x_1x_2,\ x_1x_2x_2\}$.

Moreover, let $c_w \in (\Bbbk\langle X\rangle \# \Bbbk[\Gamma])^{\chi_w}$ for all $w \in C(L)$ such that $c_w \prec_L [w]$; and let $d_u \in (\Bbbk\langle X\rangle \# \Bbbk[\Gamma])^{\chi_u^{N_u}}$ for all $u \in D(L)$ such that $d_u \prec_L [u]^{N_u}$. Then let I be the $\widehat{\Gamma}$-homogeneous ideal of $\Bbbk\langle X\rangle \# \Bbbk[\Gamma]$ generated by the following elements:

$$[w] - c_w \qquad \text{for all } w \in C(L), \tag{4.1}$$

$$[u]^{N_u} - d_u \qquad \text{for all } u \in D(L). \tag{4.2}$$

4.3 Calculation of coproducts

Let in this section $\operatorname{char}\Bbbk = 0$. For any $g \in \Gamma, \chi \in \widehat{\Gamma}$ we set

$$P_g^\chi := P_g^\chi(A) := P_{1,g}(A) \cap A^\chi = \{a \in A \mid \Delta(a) = a \otimes 1 + g \otimes a,\ ga = \chi(g)ag\}.$$

Although the following calculations are for $\Bbbk\langle X\rangle\#\Bbbk[\Gamma]$, we can use the results in any character Hopf algebra A by the canonical Hopf algebra map $\Bbbk\langle X\rangle\#\Bbbk[\Gamma] \to A$. Assume again the situation of Section 2.2.

Lemma 4.2. *Let $1 \le i < j \le \theta$ and $r \ge 1$.*

(1) *If $\operatorname{ord} q_{ii} = N$, then $x_i^N \in P_{g_i^N}^{\chi_i^N}$.*

(2) *If $q_{ij}q_{ji} = q_{ii}^{-(r-1)}$ and $r \le \operatorname{ord} q_{ii}$, then $[x_i^r x_j] \in P_{g_i^r g_j}^{\chi_i^r \chi_j}$.*

(3) *If $q_{ij}q_{ji} = q_{jj}^{-(r-1)}$ and $r \le \operatorname{ord} q_{jj}$, then $[x_i x_j^r] \in P_{g_i g_j^r}^{\chi_i \chi_j^r}$.*

Proof. (1) We have $(g_i \otimes x_i)(x_i \otimes 1) = q_{ii}(x_i \otimes 1)(g_i \otimes x_i)$ hence by Eq. (2.3) we obtain the claim. For (2) and (3) see [4, Lem. A.1]. □

Next we want to examine certain coproducts in the special case when $q_{ii} = -1$ for a $1 \le i \le \theta$. Note that in the following two Lemmata we could write more generally i and j with $1 \le i < j \le \theta$ instead of 1 and 2:

Lemma 4.3. *Let $\operatorname{ord} q_{12,12} = N$.*

(1) *If $q_{22} = -1$, we have for the quotient $(\Bbbk\langle X\rangle\#\Bbbk[\Gamma])/(x_2^2)$*

$$\begin{aligned}\Delta\big([x_1x_2]^N\big) = {}& [x_1x_2]^N \otimes 1 + g_{12}^N \otimes [x_1x_2]^N \\ & + q_{2,12}^{N-1}(1 - q_{12}q_{21})[x_1(x_1x_2)^{N-1}]g_2 \otimes x_2.\end{aligned}$$

(2) *If $q_{11} = -1$, we have for the quotient $(\Bbbk\langle X\rangle\#\Bbbk[\Gamma])/(x_1^2)$*

$$\begin{aligned}\Delta\big([x_1x_2]^N\big) = {}& [x_1x_2]^N \otimes 1 + g_{12}^N \otimes [x_1x_2]^N \\ & + q_{1,12}^{N-1}(1 - q_{12}q_{21})x_1 g_{12}^{N-1} g_2 \otimes [(x_1x_2)^{N-1}x_2].\end{aligned}$$

Proof. We calculate directly in $\Bbbk\langle X\rangle\#\Bbbk[\Gamma]$

$$\Delta([x_1x_2]) = [x_1x_2] \otimes 1 + (1 - q_{12}q_{21})x_1g_2 \otimes x_2 + g_{12} \otimes [x_1x_2].$$

For $\alpha := (1 - q_{12}q_{21})$, $q := q_{12,12}$, $U := [x_1x_2] \otimes 1$, $V := \alpha x_1 g_2 \otimes x_2$ and $W := g_{12} \otimes [x_1x_2]$ we have $WU = qUW$ and

$$\begin{aligned} VU - qUV &= \alpha q_{2,12}[x_1x_1x_2]g_2 \otimes x_2, \\ WV - qVW &= \alpha q_{12,1} x_1 g_{12} g_2 \otimes [x_1x_2x_2]. \end{aligned}$$

We further set for $r \ge 1$

$$\begin{aligned} [V] &:= V, & [VU^r] &:= \alpha q_{2,12}^r [x_1(x_1x_2)^r]g_2 \otimes x_2, \\ [W] &:= W, & [W^rV] &:= \alpha q_{1,12}^r x_1 g_{12}^r g_2 \otimes [(x_1x_2)^r x_2]. \end{aligned}$$

(1) We have $[x_1x_2x_2] = [x_1, x_2^2] = 0$ by the restricted q-Leibniz formula and $x_2^2 = 0$. Hence $WU = qUW$ and $WV = qVW$. By Eq. (2.3) we have

$$\Delta([x_1x_2]^r) = (U+V+W)^r = (U+V)^r + W^r.$$

We state for $r \geq 1$

$$(U+V)^r = U^r + \sum_{i=0}^{r-1} \tbinom{r}{i}_q U^i [VU^{(r-1)-i}],$$

from where the claim follows. This we prove by induction on r: For $r=1$ the claim is true. By induction assumption

$$\begin{aligned}
(U+V)^{r+1} &= (U+V)^r(U+V) \\
&= U^{r+1} + \sum_{i=0}^{r-1} \tbinom{r}{i}_q U^i[VU^{(r-1)-i}]U + U^rV + \sum_{i=0}^{r-1} \tbinom{r}{i}_q U^i[VU^{(r-1)-i}]V,
\end{aligned}$$

where the last sum is zero since $[VU^{(r-1)-i}]V = \ldots \otimes x_2^2 = 0$ for all $0 \leq i \leq r-1$. Further

$$\begin{aligned}
[VU^{(r-1)-i}]U &= \alpha q_{2,12}^{(r-1)-i}[x_1(x_1x_2)^{(r-1)-i}]g_2[x_1x_2] \otimes x_2 \\
&= \alpha q_{2,12}^{r-i}\big([x_1(x_1x_2)^{r-i}][x_1x_2] \\
&\qquad + q_{1,12}q^{(r-1)-i}[x_1x_2][x_1(x_1x_2)^{(r-1)-i}]\big) g_2 \otimes x_2 \\
&= [VU^{r-i}] + q^{r-i}U[VU^{(r-1)-i}].
\end{aligned}$$

Thus $(U+V)^{r+1} =$

$$\begin{aligned}
&= U^{r+1} + \sum_{i=0}^{r-1} \tbinom{r}{i}_q U^i[VU^{r-i}] + U^rV + \sum_{i=0}^{r-1} \tbinom{r}{i}_q q^{r-i} U^{i+1}[VU^{(r-1)-i}] \\
&= U^{r+1} + \sum_{i=0}^{r} \big(\tbinom{r}{i}_q + \tbinom{r}{i-1}_q q^{r+1-i}\big) U^i[VU^{r-i}],
\end{aligned}$$

by shifting the index of the second sum. Using Eq. (2.2) we get the desired formula.

(2) is proven analogously with the formula $(V+W)^r = W^r + \sum_{i=0}^{r-1} \binom{r}{i}_q [W^{(r-1)-i}V]V^i$. $\square$

A direct computation in $\Bbbk\langle X\rangle \# \Bbbk[\Gamma]$ shows that

$$\begin{aligned}
\Delta([x_1x_1x_2x_1x_2]) &= [x_1x_1x_2x_1x_2] \otimes 1 + g_1^3g_2^2 \otimes [x_1x_1x_2x_1x_2] \\
&+ \alpha[x_1x_1x_2]g_1g_2 \otimes [x_1x_2] \\
&+ (1-q_{12}q_{21})\big(q_{21}q_{22}\beta[x_1x_1x_1x_2] + \alpha[x_1x_1x_2]x_1\big) g_2 \otimes x_2 \\
&+ (1-q_{12}q_{21})(1-q_{11}q_{12}q_{21})x_1^2g_1g_2^2 \\
&\qquad \otimes \big(q_{11}q_{21}(1+q_{11}-q_{11}^3q_{12}^2q_{21}^2q_{22})[x_1x_2x_2] + \alpha x_2[x_1x_2]\big) \\
&+ q_{21}(1-q_{12}q_{21})^2(1-q_{11}q_{12}q_{21})(1-q_{11}^2q_{12}^2q_{21}^2q_{22})x_1^3g_2^2 \otimes x_2^2 \\
&+ x_1g_1^2g_2^2 \otimes \big(\gamma[x_1x_2]^2 + q_{11}^2q_{21}(1-q_{12}q_{21})[x_1x_1x_2x_2]\big),
\end{aligned}$$

with

$$\begin{aligned}
\alpha &:= (2)_{q_{11}}q_{11}q_{12}q_{21}q_{22}(1-q_{11}q_{12}q_{21}) + 1 - q_{11}^4q_{12}^3q_{21}^3q_{22}^2, \\
\beta &:= 1 - q_{11}q_{12}q_{21} - q_{11}^2q_{12}^2q_{21}^2q_{22}, \\
\gamma &:= q_{11}^2q_{21}q_{12}(1-q_{12}q_{21})(q_{22}-q_{11}) \\
&\qquad + (2)_{q_{11}}(1-q_{11}q_{12}q_{21})(1-q_{11}^3q_{12}^2q_{21}^2q_{22}).
\end{aligned}$$

Lemma 4.4. *Let* $q_{22} = -1$. *Then*

$$\alpha = (3)_{q_{12,12}}(1 - q_{11}^2 q_{12} q_{21}), \quad \beta = (3)_{q_{12,12}}, \quad \gamma = (2)_{q_{11}}(3)_{q_{12,12}}(1 - q_{11}^2 q_{12} q_{21}).$$

As a consequence we have the following:

(1) *If* $\mathrm{ord} q_{12,12} = 3$, *then*

$$\begin{aligned}
\Delta([x_1x_1x_2x_1x_2]) &= [x_1x_1x_2x_1x_2] \otimes 1 + g_1^3 g_2^2 \otimes [x_1x_1x_2x_1x_2] \\
&\quad + (1 - q_{12}q_{21})(1 - q_{11}q_{12}q_{21}) x_1^2 g_1 g_2^2 \\
&\qquad\qquad \otimes q_{11}q_{21}(1 + q_{11} - q_{11}^3 q_{12}^2 q_{21}^2 q_{22})[x_1x_2x_2] \\
&\quad + q_{21}(1 - q_{12}q_{21})^2 (1 - q_{11}q_{12}q_{21})(1 - q_{11}^2 q_{12}^2 q_{21}^2 q_{22}) x_1^3 g_2^2 \otimes x_2^2 \\
&\quad + x_1 g_1^2 g_2^2 \otimes q_{11}^2 q_{21}(1 - q_{12}q_{21})[x_1x_1x_2x_2].
\end{aligned}$$

Hence $[x_1x_1x_2x_1x_2] \in P_{g_1^3 g_2^2}^{\chi_1^3 \chi_2^2}$ *in the quotient* $(\Bbbk\langle X\rangle \# \Bbbk[\Gamma])/(x_2^2)$.

(2) *If* $q_{12}q_{21} = q_{11}^{-2}$ *and* $\mathrm{ord} q_{11} = 3$, *then*

$$\begin{aligned}
\Delta([x_1x_1x_2x_1x_2]) &= [x_1x_1x_2x_1x_2] \otimes 1 + g_1^3 g_2^2 \otimes [x_1x_1x_2x_1x_2] \\
&\quad + (1 - q_{12}q_{21}) q_{21} q_{22} \beta [x_1x_1x_1x_2] g_2 \otimes x_2 \\
&\quad + q_{21}(1 - q_{12}q_{21})^2 (1 - q_{11}q_{12}q_{21})(1 - q_{11}^2 q_{12}^2 q_{21}^2 q_{22}) x_1^3 g_2^2 \otimes x_2^2 \\
&\quad + x_1 g_1^2 g_2^2 \otimes q_{11}^2 q_{21}(1 - q_{12}q_{21})[x_1x_1x_2x_2].
\end{aligned}$$

Hence $[x_1x_1x_2x_1x_2] \in P_{g_1^3 g_2^2}^{\chi_1^3 \chi_2^2}$ *in the quotients* $(\Bbbk\langle X\rangle \# \Bbbk[\Gamma])/(x_2^2,\ [x_1x_1x_1x_2])$ *or* $(\Bbbk\langle X\rangle \# \Bbbk[\Gamma])/(x_1^3,\ [x_1x_1x_2x_2])$.

Proof. This is also a straightforward calculation using the following identities: Since $q_{22} = -1$, we have $[x_1x_2x_2] = [x_1, x_2^2]$ by the restricted q-Leibniz formula of Proposition 2.1, thus $[x_1x_1x_2x_2] = \big[x_1, [x_1x_2x_2]\big] = \big[x_1, [x_1, x_2^2]\big]$. So we see that both are zero if $x_2^2 = 0$. If $\mathrm{ord} q_{11} = 3$ analogously $[x_1x_1x_1x_2] = [x_1^3, x_2] = 0$ for $x_1^3 = 0$. $\square$

We want to state some basic combinatorics on the g_i's and χ_i's for later reference:

Lemma 4.5. *Let* $1 \le i \ne j \le \theta$, $1 < N := \mathrm{ord} q_{ii} < \infty$, *and* $r \in \mathbb{Z}$. *Then:*

(1) $\chi_i^N \ne \chi_i$.

(2) *If* $q_{jj} \ne 1$, *then* $\chi_i^N \ne \chi_j$ *or* $g_i^N \ne g_j$.

(3) *If* $\chi_i^N = \varepsilon$, *then* $q_{ji}^N = 1$. *Especially, if* $\chi_i^2 = \varepsilon$, *then* $q_{ji} = \pm 1$.

(4) *If* $q_{ij}q_{ji} = q_{ii}^{-(r-1)}$ *and* $q_{jj} \ne 1$, *then* $\chi_i^r \chi_j \ne \chi_i$.

(5) *If* $q_{ii}^r \ne 1$, *then* $\chi_i^r \chi_j \ne \chi_j$.

(6) *If* $q_{ij}q_{ji} = q_{ii}^{-(r-1)}$ *and* $\chi_i^r \chi_j = \varepsilon$, *then*

$$\begin{pmatrix} q_{ii} & q_{ij} \\ q_{ji} & q_{jj} \end{pmatrix} = \begin{pmatrix} q_{ii} & q_{ii}^{-r} \\ q_{ii} & q_{ii}^{-r} \end{pmatrix}.$$

Especially, if $q_{jj} = -1$, *then* $q_{ii}^r = -1$ *and* N *is even.*

Proof. (1) Assume $\chi_i^N = \chi_i$. Hence $q_{ii}^{N-1} = 1$, a contradiction.
(2) If $\chi_i^N = \chi_j$ and $g_i^N = g_j$, then $1 = q_{ii}^N = q_{ij}$, $q_{ji}^N = q_{jj}$, $1 = q_{ii}^N = q_{ji}$ and $q_{ij}^N = q_{jj}$. Hence $q_{jj} = q_{ij} = q_{ji} = 1$.
(3) is clear.
(4) If $\chi_i^r \chi_j = \chi_i$, then $q_{ii}^{r-1} q_{ij} = 1$, $q_{ji}^{r-1} q_{jj} = 1$. We deduce $q_{ji} = q_{jj} = 1$.
(5) If $\chi_i^r \chi_j = \chi_j$, then $q_{ii}^r = 1$.
(6) We have $q_{ii}^r q_{ij} = 1$, $q_{ji}^r q_{jj} = 1$. Now the assumption implies the claim. □

5 Lifting

We proceed as in [3, 5]: In this chapter let $\operatorname{char} \Bbbk = 0$ and A be a finite-dimensional pointed Hopf algebra with abelian group of group-like elements $G(A) = \Gamma$ and assume that the associated graded Hopf algebra with respect to the coradical filtration is

$$\operatorname{gr}(A) \cong \mathfrak{B}(V) \# \Bbbk[\Gamma],$$

where V is of diagonal type of dimension $\dim_\Bbbk V = \theta$ with basis $X = \{x_1, \ldots, x_\theta\}$. We will always identify $T(V) \cong \Bbbk\langle X \rangle$. It is $\dim_\Bbbk A = \dim_\Bbbk \operatorname{gr}(A) = \dim_\Bbbk \mathfrak{B}(V) \cdot |\Gamma|$. In particular $\mathfrak{B}(V)$ is finite-dimensional and we can associate a Cartan matrix as in Definition 1.1.

Definition 5.1. In this situation we say that A is a *lifting* of the Hopf algebra $\mathfrak{B}(V)\#\Bbbk[\Gamma]$, or simply of the Nichols algebra $\mathfrak{B}(V)$.

By [3, Lem. 5.4], we have that

$$\begin{aligned} &P_g^\varepsilon = \Bbbk(1-g) \text{ for all } g \in \Gamma, \text{ and if } \chi \neq \varepsilon, \text{ then} \\ &P_g^\chi \neq 0 \iff g = g_i,\ \chi = \chi_i \text{ for some } 1 \leq i \leq \theta. \end{aligned} \tag{5.1}$$

Thus we can choose $a_i \in P_{g_i}^{\chi_i}$ with residue class $x_i \in V\#\Bbbk[\Gamma] \cong A_1/A_0$ for $1 \leq i \leq \theta$.

Lemma 5.2. *Let $w \in C(L)$ and $u \in D(L)$.*

(1) (a) *If $q_{i,w} \neq 1$ for some $1 \leq i \leq \theta$, then $\chi_w \neq \varepsilon$.*
(b) *If $\chi_w \neq \varepsilon$ and for all $1 \leq i \leq \theta$ there are $1 \leq j \leq \theta$ such that $q_{j,w} \neq q_{ji}$ or $q_{w,j} \neq q_{ij}$, then*

$$P_{g_w}^{\chi_w} = 0.$$

(2) *Let* $\operatorname{ord} q_{u,u} = N_u < \infty$.
(a) *If $q_{i,u}^{N_u} \neq 1$ for some $1 \leq i \leq \theta$, then $\chi_u^{N_u} \neq \varepsilon$.*
(b) *If $\chi_u^{N_u} \neq \varepsilon$ and for all $1 \leq i \leq \theta$ there are $1 \leq j \leq \theta$ such that $q_{j,u}^{N_u} \neq q_{ji}$ or $q_{u,j}^{N_u} \neq q_{ij}$, then*

$$P_{g_u^{N_u}}^{\chi_u^{N_u}} = 0.$$

Proof. (1a) If $\chi_w = \varepsilon$, then $q_{i,w} = 1$ for all $1 \leq i \leq \theta$.
(1b) Let $\chi_w \neq \varepsilon$ and $P_{g_w}^{\chi_w} \neq 0$, then $\chi_w = \chi_i$ and $g_w = g_i$ for some i by Eq. (5.1). Hence $q_{j,w} = q_{ji}$ and $q_{w,j} = q_{ij}$ for all $1 \leq j \leq \theta$.
(2a) If $\chi_u^{N_u} = \varepsilon$, then $q_{i,u}^{N_u} = 1$ for all $1 \leq i \leq \theta$.
(2b) Let $\chi_u^{N_u} \neq \varepsilon$ and $P_{g_u^{N_u}}^{\chi_u^{N_u}} \neq 0$, then $\chi_u^{N_u} = \chi_i$ and $g_u^{N_u} = g_i$ for some i. Thus $q_{j,u}^{N_u} = q_{ji}$ and $q_{u,j}^{N_u} = q_{ij}$ for all $1 \leq j \leq \theta$. □

This and Eq. (5.1) motivate the following:

Definition 5.3. Let $L \subset \mathcal{L}$. Then we define coefficients $\mu_u \in \Bbbk$ for all $u \in D(L)$, and $\lambda_w \in \Bbbk$ for all $w \in C(L)$ by

$$\mu_u = 0, \text{ if } g_u^{N_u} = 1 \text{ or } \chi_u^{N_u} \neq \varepsilon, \qquad \lambda_w = 0, \text{ if } g_w = 1 \text{ or } \chi_w \neq \varepsilon,$$

and otherwise they can be chosen arbitrarily.

5.1 General lifting procedure

Suppose we know the PBW basis $[L]$ of $\mathfrak{B}(V)$, then a lifting A has the same PBW basis $[L]$; see [29, Prop. 47]. Hence we know by Section 4 the structure of the ideal I such that

$$A \cong (\Bbbk\langle X\rangle \# \Bbbk[\Gamma])/I.$$

Let us order the relations Eqs. (4.1) and (4.2) of I, namely the two types

$$[w] - c_w \text{ for } w \in C(L) \quad \text{and} \quad [u]^{N_u} - d_u \text{ for } u \in D(L),$$

with respect to $\prec$ by the leading super word $[w]$ resp. $[u]^{N_u}$. Yet we don't know the $c_w, d_u \in \Bbbk\langle X\rangle \# \Bbbk[\Gamma]$ explicitly; our general procedure to compute these elements is the following, stated inductively on $\prec$:

- Suppose we know all relations $\prec$-smaller than $[w]$ resp. $[u]^{N_u}$.
- Then we determine a counterterm r_w resp. $s_u \in \Bbbk\langle X\rangle \# \Bbbk[\Gamma]$ such that

  $$[w] - r_w \in P_{g_w}^{\chi_w} \quad \text{resp.} \quad [u]^{N_u} - s_u \in P_{g_u^{N_u}}^{\chi_u^{N_u}}$$

  modulo the relations $\prec$-smaller than $[w]$ resp. $[u]^{N_u}$; we conjecture that we can do this in general (see below).
  Further if $\chi_w \neq \chi_i$ or $g_w \neq g_i$ resp. $\chi_u^{N_u} \neq \chi_i$ or $g_u^{N_u} \neq g_i$ for all $1 \leq i \leq \theta$, then by Eq. (5.1) we get

  $$\boxed{c_w = r_w + \lambda_w(1 - g_w) \quad \text{resp.} \quad d_u = s_u + \mu_u(1 - g_u^{N_u}).} \tag{5.2}$$

In order to formulate our conjecture, we define the following ideal: For any super word $U \in [\mathcal{L}]^{(\mathbb{N})}$ let I_U denote the ideal of $\Bbbk\langle X\rangle \# \Bbbk[\Gamma]$ generated by the elements

$$\begin{aligned} &[w] - c_w && \text{for all } w \in C(L) \text{ and } [w] \prec U, \\ &[u]^{N_u} - d_u && \text{for all } u \in D(L) \text{ and } [u]^{N_u} \prec U. \end{aligned}$$

Note that $I_U \subset I$.

Conjecture 5.4. *For all $w \in C(L)$ resp. for all $u \in D(L)$ there are $r_w \in (\Bbbk\langle X\rangle \# \Bbbk[\Gamma])^{\chi_w}$ resp. $s_u \in (\Bbbk\langle X\rangle \# \Bbbk[\Gamma])^{\chi_u^{N_u}}$ with $r_w \prec_L [w]$ resp. $s_u \prec_L [u]^{N_u}$ such that $[w] - r_w$ resp. $[u]^{N_u} - s_u$ is skew-primitive modulo the relations $\prec$-smaller than $[w]$ resp. $[u]^{N_u}$, i.e.,*

$$\begin{aligned} &\Delta([w] - r_w) - ([w] - r_w) \otimes 1 - g_w \otimes ([w] - r_w) \\ &\qquad \in \Bbbk\langle X\rangle \# \Bbbk[\Gamma] \otimes I_{[w]} + I_{[w]} \otimes \Bbbk\langle X\rangle \# \Bbbk[\Gamma], \\ &\Delta([u]^{N_u} - s_u) - ([u]^{N_u} - s_u) \otimes 1 - g_u^{N_u} \otimes ([u]^{N_u} - s_u) \\ &\qquad \in \Bbbk\langle X\rangle \# \Bbbk[\Gamma] \otimes I_{[u]^{N_u}} + I_{[u]^{N_u}} \otimes \Bbbk\langle X\rangle \# \Bbbk[\Gamma]. \end{aligned}$$

Remark 5.5.

1. If the conjecture is true, then one could investigate from the list of braidings in [19] where a free paramter λ_w resp. μ_u occurs in the lifting, without knowing r_w resp. s_u explicitly.

2. To determine the generators of the ideal I explicitly, i.e., to find r_w resp. s_u, it is crucial to know which relations of I are redundant. The redundant relations are detected as described in [23].

3. In general r_w resp. s_u is not necessarily in $\Bbbk[\Gamma]$, like it was the case in [7]; see Lemma 5.6 (2b),(3b) below or the liftings in the following sections.

At first we lift the root vector relations of $x_1, \dots, x_\theta$ and the Serre relations in general. Note that for these relations our Conjecture 5.4 is true. We denote the images of $[x_i^r x_j], [x_i x_j^r] \in \Bbbk\langle X\rangle$ $(r \geq 1)$ of the algebra map $\Bbbk\langle X\rangle \to A, \quad x_i \mapsto a_i$ by $[a_i^r a_j], [a_i a_j^r]$:

Lemma 5.6. *Let A be a lifting of $\mathfrak{B}(V)$ with braiding matrix (q_{ij}) and Cartan matrix (a_{ij}). Further let $1 \leq i < j \leq \theta$ and $N_i := \mathrm{ord} q_{ii}$. We may assume $q_{ii} \neq 1$ for all $1 \leq i \leq \theta$.*

(1) *We have*

$$a_i^{N_i} = \mu_i(1 - g_i^{N_i}).$$

Moreover, if $q_{ji}^{N_i} \neq 1$, then $a_i^{N_i} = 0$. Especially, if $q_{ii} = -1$ and $q_{ji} \neq \pm 1$, then $a_i^2 = 0$.

(2) (a) *If $q_{ij}q_{ji} = q_{ii}^{a_{ij}}$, $N_i > 1 - a_{ij}$, then*

$$\left[a_i^{1-a_{ij}} a_j\right] = \lambda_{i^{1-a_{ij}}j}(1 - g_i^{1-a_{ij}} g_j).$$

Moreover, if $\begin{pmatrix} q_{ii} & q_{ij} \\ q_{ji} & q_{jj} \end{pmatrix} \neq \begin{pmatrix} q_{ii} & q_{ii}^{-(1-a_{ij})} \\ q_{ii} & q_{ii}^{-(1-a_{ij})} \end{pmatrix}$, then $[a_i^{1-a_{ij}} a_j] = 0$; in particular the latter claim holds if $q_{jj} = -1$ and $q_{ii}^{1-a_{ij}} \neq -1$ (e.g., N_i is odd).

(b) *If $N_i = 1 - a_{ij}$, then*

$$\left[a_i^{N_i} a_j\right] = \mu_i(1 - q_{ij}^{N_i})a_j.$$

(3) (a) *If $q_{ij}q_{ji} = q_{jj}^{a_{ji}}$, $N_j > 1 - a_{ji}$, then*

$$\left[a_i a_j^{1-a_{ji}}\right] = \lambda_{ij^{1-a_{ji}}}(1 - g_i g_j^{1-a_{ji}}).$$

Moreover, if $\begin{pmatrix} q_{ii} & q_{ij} \\ q_{ji} & q_{jj} \end{pmatrix} \neq \begin{pmatrix} q_{jj}^{-(1-a_{ji})} & q_{jj} \\ q_{jj}^{-(1-a_{ji})} & q_{jj} \end{pmatrix}$, then $[a_i a_j^{1-a_{ji}}] = 0$; in particular the latter claim holds if $q_{ii} = -1$ and $q_{jj}^{1-a_{ji}} \neq -1$ (e.g., N_j is odd).

(b) *If $N_j = 1 - a_{ji}$, then*

$$\left[a_i a_j^{N_j}\right] = \mu_j(q_{ji}^{N_j} - 1)a_i g_j^{N_j}.$$

Proof. (1) This is a consequence of Lemma 4.5(1)-(3) and Eq. (5.1).

(2a) and (3a) follow from Lemma 4.5(4)-(6) and Eq. (5.1).

(2b) and (3b) follow from the restricted q-Leibniz formula of Proposition 2.1 and (1) above: For example

$$\left[a_i a_j^{N_j}\right] = \left[a_i, a_j^{N_j}\right]_{q_{ij}^{N_j}} = \left[a_i, \mu_j(1 - g_j^{N_j})\right]_{q_{ij}^{N_j}} = \mu_j\Big((1 - q_{ij}^{N_j})a_i - (1 - q_{ij}^{N_j} q_{ji}^{N_j})a_i g_j^{N_j}\Big).$$

Now either $\mu_j = 0$ or $q_{ij}^{N_j} = 1$ by (1), from where the claim follows. $\square$

From now on let $\theta = 2$, i.e., $\mathfrak{B}(V)$ is of rank 2.

5.2 Lifting of $\mathfrak{B}(V)$ with Cartan matrix $A_1 \times A_1$

Let $\mathfrak{B}(V)$ be a finite-dimensional Nichols algebras with Cartan matrix $(a_{ij}) = \left(\begin{smallmatrix} 2 & 0 \\ 0 & 2 \end{smallmatrix}\right)$ of type $A_1 \times A_1$, i.e., the braiding matrix (q_{ij}) fulfills

$$q_{12}q_{21} = 1,$$

since we may suppose that $\mathrm{ord} q_{ii} \geq 2$ [15, Sect. 2], especially $q_{ii} \neq 1$. The Dynkin diagram is $\overset{q}{\circ} \qquad \overset{r}{\circ}$ with $q := q_{11}$ and $r := q_{22}$. Then the Nichols algebra is given by

$$\mathfrak{B}(V) = T(V)/\big([x_1x_2],\, x_1^{N_1},\, x_2^{N_2}\big)$$

with basis $\{x_2^{r_2}x_1^{r_1} \mid 0 \leq r_i < N_i\}$ where $N_i = \mathrm{ord} q_{ii} \geq 2$ [18]. It is well-known [3] that any lifting A is of the form

$$\begin{aligned} A \cong (T(V)\#\Bbbk[\Gamma])/\big(\quad & [x_1x_2] - \lambda_{12}(1 - g_{12}),\ x_1^{N_1} - \mu_1(1 - g_1^{N_1}), \\ & \qquad\qquad\qquad\quad\ x_2^{N_2} - \mu_2(1 - g_2^{N_2}) \quad \big) \end{aligned}$$

with basis $\{x_2^{r_2}x_1^{r_1}g \mid 0 \leq r_i < N_i, g \in \Gamma\}$ and $\dim_\Bbbk A = N_1N_2 \cdot |\Gamma|$; the statement for the basis is proven in [23].

5.3 Lifting of $\mathfrak{B}(V)$ with Cartan matrix A_2

Let $\mathfrak{B}(V)$ be a Nichols algebras with Cartan matrix $(a_{ij}) = \left(\begin{smallmatrix} 2 & -1 \\ -1 & 2 \end{smallmatrix}\right)$ of type A_2, i.e., the braiding matrix (q_{ij}) fulfills

$$q_{12}q_{21} = q_{11}^{-1} \text{ or } q_{11} = -1, \text{ and } q_{12}q_{21} = q_{22}^{-1} \text{ or } q_{22} = -1.$$

The Nichols algebras are given explicitly in [17]. As mentioned above, it is crucial to know the redundant relations for the computation of the liftings. Therefore we give the ideals without redundant relations which are detected in [23]:

Proposition 5.7 (Nichols algebras with Cartan matrix A_2)**.** *The finite-dimensional Nichols algebras $\mathfrak{B}(V)$ with Cartan matrix of type A_2 are exactly the following:*

(1) $\overset{q}{\circ}\overset{q^{-1}}{\text{———}}\overset{q}{\circ}$ *(Cartan type A_2). Let $q_{12}q_{21} = q_{11}^{-1} = q_{22}^{-1}$.*

(a) *If $q_{11} = -1$, then*

$$\mathfrak{B}(V) = T(V)/\big(x_1^2,\ [x_1x_2]^2,\ x_2^2\big)$$

with basis $\{x_2^{r_2}[x_1x_2]^{r_{12}}x_1^{r_1} \mid 0 \leq r_2, r_{12}, r_1 < 2\}$ and $\dim_\Bbbk \mathfrak{B}(V) = 2^3 = 8$.

(b) *If $N := \mathrm{ord} q_{11} \geq 3$, then*

$$\mathfrak{B}(V) = T(V)/\big([x_1x_1x_2],\ [x_1x_2x_2],\ x_1^N,\ [x_1x_2]^N,\ x_2^N\big)$$

with basis $\{x_2^{r_2}[x_1x_2]^{r_{12}}x_1^{r_1} \mid 0 \leq r_2, r_{12}, r_1 < N\}$ and $\dim_\Bbbk \mathfrak{B}(V) = N^3$.

(2) $\overset{q}{\circ}\overset{q^{-1}}{\text{———}}\overset{-1}{\circ}$. *If $q_{12}q_{21} = q_{11}^{-1}$, $N := \mathrm{ord} q_{11} \geq 3$, $q_{22} = -1$, then*

$$\mathfrak{B}(V) = T(V)/\big([x_1x_1x_2],\ x_1^N,\ x_2^2\big)$$

with basis $\{x_2^{r_2}[x_1x_2]^{r_{12}}x_1^{r_1} \mid 0 \leq r_1 < N,\, 0 \leq r_2, r_{12} < 2\}$ and $\dim_\Bbbk \mathfrak{B}(V) = 4N$.

(3) $\overset{-1}{\circ}\overset{q^{-1}}{\text{——}}\overset{q}{\circ}$. *If* $q_{11}=-1$, $q_{12}q_{21}=q_{22}^{-1}$, $N:=\mathrm{ord}q_{22}\geq 3$, *then*

$$\mathfrak{B}(V)=T(V)/\big([x_1x_2x_2],\ x_1^2,\ x_2^N\big)$$

with basis $\{x_2^{r_2}[x_1x_2]^{r_{12}}x_1^{r_1} \mid 0\leq r_2<N,\, 0\leq r_1,r_{12}<2\}$ *and* $\dim_{\Bbbk}\mathfrak{B}(V)=4N$.

(4) $\overset{-1}{\circ}\overset{q}{\text{——}}\overset{-1}{\circ}$. *If* $q_{11}=q_{22}=-1$, $N:=\mathrm{ord}q_{12}q_{21}\geq 3$, *then*

$$\mathfrak{B}(V)=T(V)/\big(x_1^2,\ [x_1x_2]^N,\ x_2^2\big)$$

with basis $\{x_2^{r_2}[x_1x_2]^{r_{12}}x_1^{r_1} \mid 0\leq r_2,r_1<2,\, 0\leq r_{12}<N\}$ *and* $\dim_{\Bbbk}\mathfrak{B}(V)=4N$.

Remark 5.8. The Nichols algebras of Proposition 5.7 all have the PBW basis $[L]=\{x_1,[x_1x_2],x_2\}$, and (1) resp. (2)-(4) form the standard Weyl equivalence class of row 2 resp. 3 in [15, 18, Figure 1], where the latter is not of Cartan type. They build up the tree type T_2 of [17].

Theorem 5.9 (Liftings of $\mathfrak{B}(V)$ with Cartan matrix A_2)**.** *For any lifting A of $\mathfrak{B}(V)$ as in Proposition 5.7, we have*

$$A\cong (T(V)\#\Bbbk[\Gamma])/I,$$

where I is specified as follows:

(1) $\overset{q}{\circ}\overset{q^{-1}}{\text{——}}\overset{q}{\circ}$ *(Cartan type A_2). Let* $q_{12}q_{21}=q_{11}^{-1}=q_{22}^{-1}$.

(a) *If* $q_{11}=-1$, *then I is generated by*

$$\begin{aligned} x_1^2-\mu_1(1-g_1^2),\\ [x_1x_2]^2-4\mu_1q_{21}x_2^2-\mu_{12}(1-g_{12}^2),\\ x_2^2-\mu_2(1-g_2^2). \end{aligned}$$

A basis is $\{x_2^{r_2}[x_1x_2]^{r_{12}}x_1^{r_1}g \mid 0\leq r_2,r_{12},r_1<2,\ g\in\Gamma\}$ *and* $\dim_{\Bbbk}A=2^3\cdot|\Gamma|=8\cdot|\Gamma|$.

(b) *If* $\mathrm{ord}q_{11}=3$, *then I is generated by, see* [10],

$$\begin{aligned} &[x_1x_1x_2]-\lambda_{112}(1-g_{112}), && x_1^3-\mu_1(1-g_1^3),\\ &[x_1x_2x_2]-\lambda_{122}(1-g_{122}), && [x_1x_2]^3+(1-q_{11})q_{11}\lambda_{112}[x_1x_2x_2]\\ & && \quad -\mu_1(1-q_{11})^3x_2^3-\mu_{12}(1-g_{12}^3),\\ & && x_2^3-\mu_2(1-g_2^3). \end{aligned}$$

A basis is $\{x_2^{r_2}[x_1x_2]^{r_{12}}x_1^{r_1}g \mid 0\leq r_2,r_{12},r_1<3,\ g\in\Gamma\}$ *and* $\dim_{\Bbbk}A=3^3\cdot|\Gamma|=27\cdot|\Gamma|$.

(c) *If* $N:=\mathrm{ord}q_{11}\geq 4$, *then I is generated by, see* [5],

$$\begin{aligned} &[x_1x_1x_2], && x_1^N-\mu_1(1-g_1^N),\\ &[x_1x_2x_2], && [x_1x_2]^N-\mu_1(q_{11}-1)^Nq_{21}^{\frac{N(N-1)}{2}}x_2^N-\mu_{12}(1-g_{12}^N),\\ & && x_2^N-\mu_2(1-g_2^N). \end{aligned}$$

A basis is $\{x_2^{r_2}[x_1x_2]^{r_{12}}x_1^{r_1}g \mid 0\leq r_2,r_{12},r_1<N,\ g\in\Gamma\}$ *and* $\dim_{\Bbbk}A=N^3\cdot|\Gamma|$.

(2) 

Let $q_{12}q_{21} = q_{11}^{-1}$, $q_{22} = -1$.

(a) *If* $4 \neq N := \mathrm{ord} q_{11} \geq 3$, *then* I *is generated by*

$$[x_1x_1x_2], \qquad x_1^N - \mu_1(1-g_1^N), \quad x_2^2 - \mu_2(1-g_2^2).$$

A basis is $\{x_2^{r_2}[x_1x_2]^{r_{12}}x_1^{r_1}g \mid 0 \leq r_1 < N, 0 \leq r_2, r_{12} < 2, g \in \Gamma\}$ *and* $\dim_{\Bbbk} A = 2^2N \cdot |\Gamma| = 4N \cdot |\Gamma|$.

(b) *If* $\mathrm{ord} q_{11} = 4$, *then* I *is generated by*

$$[x_1x_1x_2] - \lambda_{112}(1-g_{112}), \qquad x_1^4 - \mu_1(1-g_1^4), \quad x_2^2 - \mu_2(1-g_2^2).$$

A basis is $\{x_2^{r_2}[x_1x_2]^{r_{12}}x_1^{r_1}g \mid 0 \leq r_1 < 4, 0 \leq r_2, r_{12} < 2, g \in \Gamma\}$ *and* $\dim_{\Bbbk} A = 2^2 4 \cdot |\Gamma| = 16 \cdot |\Gamma|$.

(3) 

Let $q_{11} = -1$, $q_{12}q_{21} = q_{22}^{-1}$.

(a) *If* $4 \neq N := \mathrm{ord} q_{22} \geq 3$, *then* I *is generated by*

$$[x_1x_2x_2], \qquad x_1^2 - \mu_1(1-g_1^2), \quad x_2^N - \mu_2(1-g_2^N).$$

A basis is $\{x_2^{r_2}[x_1x_2]^{r_{12}}x_1^{r_1}g \mid 0 \leq r_2 < N, 0 \leq r_1, r_{12} < 2, g \in \Gamma\}$ *and* $\dim_{\Bbbk} A = 2^2N \cdot |\Gamma| = 4N \cdot |\Gamma|$.

(b) *If* $\mathrm{ord} q_{22} = 4$, *then* I *is generated by*

$$[x_1x_2x_2] - \lambda_{122}(1-g_{122}), \qquad x_1^2 - \mu_1(1-g_1^2), \quad x_2^4 - \mu_2(1-g_2^4).$$

A basis is $\{x_2^{r_2}[x_1x_2]^{r_{12}}x_1^{r_1}g \mid 0 \leq r_2 < 4, 0 \leq r_1, r_{12} < 2, g \in \Gamma\}$ *and* $\dim_{\Bbbk} A = 2^2 4 \cdot |\Gamma| = 16 \cdot |\Gamma|$.

(4) 

Let $q_{11} = q_{22} = -1$ *and* $N := \mathrm{ord} q_{12}q_{21} \geq 3$.

(a) *If* $q_{12} \neq \pm 1$, *then* I *is generated by*

$$x_1^2 - \mu_1(1-g_1^2), \quad [x_1x_2]^N - \mu_{12}(1-g_{12}^N), \quad x_2^2.$$

(b) *If* $q_{12} = \pm 1$, *then* I *is generated by*

$$x_1^2, \quad [x_1x_2]^N - \mu_{12}(1-g_{12}^N), \quad x_2^2 - \mu_2(1-g_2^2).$$

In both cases a basis is $\{x_2^{r_2}[x_1x_2]^{r_{12}}x_1^{r_1}g \mid 0 \leq r_2, r_1 < 2, 0 \leq r_{12} < N, g \in \Gamma\}$ *and* $\dim_{\Bbbk} A = 2^2N \cdot |\Gamma| = 4N \cdot |\Gamma|$.

Proof. At first we show that in each case $(T(V)\#\Bbbk[\Gamma])/I$ is a pointed Hopf algebra with coradical $\Bbbk[\Gamma]$ and claimed basis and dimension such that $\operatorname{gr}((T(V)\#\Bbbk[\Gamma])/I) \cong \mathfrak{B}(V)\#\Bbbk[\Gamma]$. Then we show that a lifting A is necessarily of this form.

- $(T(V)\#\Bbbk[\Gamma])/I$ is a Hopf algebra: We show that in every case I is generated by skew-primitive elements, thus I is a Hopf ideal. The elements $x_i^{N_i} - \mu_i(1 - g_i^{N_i})$ and $[x_1x_1x_2] - \lambda_{112}(1 - g_{112})$ are skew-primitive if $q_{12}q_{21} = q_{11}^{-1}$ by Lemma 4.2. So we have a Hopf ideal in (2) and (3).

For the elements $[x_1x_2]^{N_{12}} - d_{12}$ we argue as follows: In (1a) we directly calculate that $[x_1x_2]^2 - 4\mu_1 q_{21} x_2^2 \in P^{\chi_{12}^2}_{g_{12}^2}$. (1b),(1c) is treated in [10, 5].

For (4a): By induction on N (the induction basis $N = 2$ is Lemma 5.6(2b))

$$[x_1(x_1x_2)^{N-1}] = \mu_1\Big(\prod_{i=0}^{N-2}(1 - q_{12}^{i+2}q_{21}^{i})\Big)x_2[x_1x_2]^{N-2}.$$

Further $q_{12,12} = q_{12}q_{21}$ is of order N and $q_{21}^2 = 1$ (or $\mu_1 = 0$), we have $q_{12}^N q_{21}^{N-2} = (q_{12}q_{21})^r = 1$ and thus $[x_1(x_1x_2)^{N-1}] = 0$. Hence $[x_1x_2]^N$ is skew-primitive by Lemma 4.3(1).

(4b) works in a similar way because of $q_{12}^2 = 1$: Again by induction (the induction basis $N = 2$ is Lemma 5.6(3b))

$$[(x_1x_2)^{N-1}x_2] = \mu_2\Big(\prod_{i=0}^{N-2}(1 - q_{12}^{i+2}q_{21}^{i+2})\Big)[x_1x_2]^{N-2}x_1g_2^2,$$

which is 0 since $(q_{12}q_{21})^N = q_{12,12}^N = 1$. Now $[x_1x_2]^N$ is skew-primitive by Lemma 4.3(2).

- The statement on the basis and dimension of $(T(V)\#\Bbbk[\Gamma])/I$ is proven in [23].
- The algebra $\Bbbk[\Gamma]$ embeds in $(T(V)\#\Bbbk[\Gamma])/I$ and the coradical of the latter is $((T(V)\#\Bbbk[\Gamma])/I)_0 = \Bbbk[\Gamma]$ [26, Lem. 5.5.1], so $(T(V)\#\Bbbk[\Gamma])/I$ is pointed.
- We consider the Hopf algebra map

$$T(V)\#\Bbbk[\Gamma] \to \operatorname{gr}((T(V)\#\Bbbk[\Gamma])/I)$$

which maps x_i onto the residue class of x_i in the homogeneous component of degree 1, namely $((T(V)\#\Bbbk[\Gamma])/I)_1/\Bbbk[\Gamma]$. It is surjective, since $(T(V)\#\Bbbk[\Gamma])/I$ is generated as an algebra by x_1, x_2 and Γ. Further it factorizes to

$$\mathfrak{B}(V)\#\Bbbk[\Gamma] \xrightarrow{\sim} \operatorname{gr}((T(V)\#\Bbbk[\Gamma])/I).$$

This is a direct argument looking at the coradical filtration as in [3, Cor. 5.3]: all equations of I are of the form $[w] - c_w$, $[u]^{N_u} - d_u$ with $c_w, d_u \in \Bbbk[\Gamma] = ((T(V)\#\Bbbk[\Gamma])/I)_0$, hence $[w] = 0$, $[u]^{N_u} = 0$ in $\operatorname{gr}((T(V)\#\Bbbk[\Gamma])/I)$. The latter surjective Hopf algebra map must be an isomorphism because the dimensions coincide.

- The other way round, let A be a lifting of $\mathfrak{B}(V)$ with $a_i \in P^{\chi_i}_{g_i}$ as in the beginning of this chapter. We consider the Hopf algebra map

$$T(V)\#\Bbbk[\Gamma] \to A$$

which takes x_i to a_i and g to g. It is surjective since A is generated by a_1, a_2 and Γ [3, Lem. 2.2]. We have to check whether this map factorizes to

$$(T(V)\#\Bbbk[\Gamma])/I \xrightarrow{\sim} A.$$

Then we are done since the dimension implies that this is an isomorphism.

But this means we have to check that the relations of I hold in A: By Lemma 5.6 the relations concerning the elements $a_i^{N_i}$, $[a_1a_1a_2]$ and $[a_1a_2a_2]$ are of the right form. We are left to check those for $[a_1a_2]^{N_{12}}$, which appear in (1) and (4):

In (1a) we have $[a_1a_2]^2 - 4\mu_1 q_{21} a_2^2 \in P_{g_{12}^2}^{\chi_{12}^2}$ like before. Now since $q_{1,12}^2 = q_{12}^2 \neq -1 = q_{11}$ or $q_{12,2}^2 = q_{12}^2 \neq q_{12}$, and $q_{1,12}^2 = q_{12}^2 \neq q_{12}$ or $q_{12,2}^2 = q_{12}^2 \neq -1 = q_{22}$ (otherwise we get the contradiction $q_{12} = 1$ and $q_{12}^2 = -1$), we have $[a_1a_2]^2 = 4\mu_1 q_{21} a_2^2 + \mu_{12}(1 - g_{12}^2)$ by Lemma 5.2(2). (1b),(1c) work in the same way; see [10, 5]. For (4): As shown before $[a_1a_2]^N \in P_{g_{12}^N}^{\chi_{12}^N}$. Again we deduce from Lemma 5.2(2) that $[a_1a_2]^N = \mu_{12}(1 - g_{12}^N)$. $\square$

Remark 5.10. The Conjecture 5.4 is true in the situation of Theorem 5.9: the r_w of the non-redundant relations $[w] - c_w$ are 0 ($r_{112} = r_{122} = 0$ if the Serre relations are not redundant) and $s_{12} \in \Bbbk[\Gamma]$ in (1), otherwise $s_u = 0$ if $[u]^{N_u} - d_u$ is not redundant.

5.4 Lifting of $\mathfrak{B}(V)$ with Cartan matrix B_2

In this section we lift some of the Nichols algebras of standard type with associated Cartan matrix $\left(\begin{smallmatrix} 2 & -3 \\ -1 & 2 \end{smallmatrix}\right)$ of type B_2 (in the next Section also of non-standard type B_2). At first we recall the Nichols algebras (see [17]), but again we give the ideals without redundant relations [23]:

Proposition 5.11 (Nichols algebras with Cartan matrix B_2)**.** *The following finite-dimensional Nichols algebras $\mathfrak{B}(V)$ of standard type with braiding matrix (q_{ij}) and Cartan matrix of type B_2 are represented as follows:*

(1) 

(Cartan type B_2). Let $q_{12}q_{21} = q_{11}^{-2} = q_{22}^{-1}$ and $N := \operatorname{ord} q_{11}$.

(a) *If $N = 3$, then*

$$\mathfrak{B}(V) = T(V)/\big([x_1x_2x_2],\ x_1^3,\ [x_1x_1x_2]^3,\ [x_1x_2]^3,\ x_2^3\big)$$

with basis $\big\{x_2^{r_2}[x_1x_2]^{r_{12}}[x_1x_1x_2]^{r_{112}}x_1^{r_1} \mid 0 \leq r_1, r_{12}, r_{112}, r_2 < 3\big\}$ and $\dim_\Bbbk \mathfrak{B}(V) = 3^4 = 81$.

(b) *If $N = 4$, then*

$$\mathfrak{B}(V) = T(V)/\big([x_1x_1x_1x_2],\ x_1^4,\ [x_1x_1x_2]^2,\ [x_1x_2]^4,\ x_2^2\big)$$

with basis $\big\{x_2^{r_2}[x_1x_2]^{r_{12}}[x_1x_1x_2]^{r_{112}}x_1^{r_1} \mid 0 \leq r_1, r_{12} < 4,\ 0 \leq r_2, r_{112} < 2\big\}$ and $\dim_\Bbbk \mathfrak{B}(V) = 2^2 \cdot 4^2 = 64$.

(c) *If $N \geq 5$ is odd, then*

$$\mathfrak{B}(V) = T(V)/\big([x_1x_1x_1x_2],\ [x_1x_2x_2],\ x_1^N,\ [x_1x_1x_2]^N,\ [x_1x_2]^N,\ x_2^N\big)$$

with basis $\big\{x_2^{r_2}[x_1x_2]^{r_{12}}[x_1x_1x_2]^{r_{112}}x_1^{r_1} \mid 0 \leq r_1, r_{12}, r_{112}, r_2 < N\big\}$ and $\dim_\Bbbk \mathfrak{B}(V) = N^4$.

(d) *If $N \geq 6$ is even, then*

$$\mathfrak{B}(V) = T(V)/\big([x_1x_1x_1x_2],\ [x_1x_2x_2],\ x_1^N,\ [x_1x_1x_2]^{\frac{N}{2}},\ [x_1x_2]^N,\ x_2^{\frac{N}{2}}\big)$$

with basis $\big\{x_2^{r_2}[x_1x_2]^{r_{12}}[x_1x_1x_2]^{r_{112}}x_1^{r_1} \mid 0 \leq r_1, r_{12} < N,\ 0 \leq r_2, r_{112} < \frac{N}{2}\big\}$ and $\dim_\Bbbk \mathfrak{B}(V) = \frac{N^4}{4}$.

(2) 

Let $q_{12}q_{21} = q_{11}^{-2}$, $q_{22} = -1$ *and* $N := \mathrm{ord} q_{11}$.

(a) *If* $N = 3$, *then*

$$\mathfrak{B}(V) = T(V)/\big([x_1x_1x_2x_1x_2],\ x_1^3,\ [x_1x_2]^6,\ x_2^2\big)$$

with basis $\big\{x_2^{r_2}[x_1x_2]^{r_{12}}[x_1x_1x_2]^{r_{112}}x_1^{r_1} \mid 0 \le r_1 < 3,\ 0 \le r_{12} < 6,\ 0 \le r_2, r_{112} < 2\ \big\}$ *and* $\dim_{\Bbbk} \mathfrak{B}(V) = 72$.

(b) *If* $N \ge 5$ *(*$N = 4$ *is* (1b)*), then for* $N' := \mathrm{ord}(-q_{11}^{-1})$

$$\mathfrak{B}(V) = T(V)/\big([x_1x_1x_1x_2],\ x_1^N,\ [x_1x_2]^{N'},\ x_2^2\big)$$

with basis $\big\{x_2^{r_2}[x_1x_2]^{r_{12}}[x_1x_1x_2]^{r_{112}}x_1^{r_1} \mid 0 \le r_1 < N,\ 0 \le r_{12} < N',\ 0 \le r_2, r_{112} < 2\big\}$ *and* $\dim_{\Bbbk} \mathfrak{B}(V) = 4NN'$.

(3) 

Let $\mathrm{ord} q_{11} = 3$, $q_{12}q_{21} = q_{22}^{-1}$ *and* $N := \mathrm{ord} q_{22}$.

(a) *If* $N = 2$, *then*

$$\mathfrak{B}(V) = T(V)/\big([x_1x_1x_2x_1x_2],\ x_1^3,\ [x_1x_1x_2]^6,\ x_2^2\big)$$

with basis $\big\{x_2^{r_2}[x_1x_2]^{r_{12}}[x_1x_1x_2]^{r_{112}}x_1^{r_1} \mid 0 \le r_1, r_{12} < 3,\ 0 \le r_2 < 2,\ 0 \le r_{112} < 6\big\}$ *and* $\dim_{\Bbbk} \mathfrak{B}(V) = 108$.

(b) *If* $N \ge 4$ *(*$N = 3$ *is* (1) *or Proposition 5.7*(1)*), then for* $N' := \mathrm{ord} q_{11}q_{22}^{-1}$

$$\mathfrak{B}(V) = T(V)/\big([x_1x_2x_2],\ x_1^3,\ [x_1x_1x_2]^{N'},\ x_2^N\big)$$

with basis $\big\{x_2^{r_2}[x_1x_2]^{r_{12}}[x_1x_1x_2]^{r_{112}}x_1^{r_1} \mid 0 \le r_1, r_{12} < 3,\ 0 \le r_2 < N,\ 0 \le r_{112} < N'\big\}$ *and* $\dim_{\Bbbk} \mathfrak{B}(V) = 9NN'$.

(4) 

Let $\mathrm{ord} q_{11} = 3$, $q_{12}q_{21} = -q_{11}$, $q_{22} = -1$, *then*

$$\mathfrak{B}(V) = T(V)/\big([x_1x_1x_2x_1x_2],\ x_1^3,\ x_2^2\big)$$

with basis $\big\{x_2^{r_2}[x_1x_2]^{r_{12}}[x_1x_1x_2]^{r_{112}}x_1^{r_1} \mid 0 \le r_1, r_{12} < 3,\ 0 \le r_2, r_{112} < 2\big\}$ *and* $\dim_{\Bbbk} A = 36$.

Remark 5.12. The Nichols algebras of Proposition 5.11 all have the PBW basis $[L] = \{x_2, [x_1x_2], [x_1x_1x_2], x_1\}$, and (1)-(4) form the standard Weyl equivalence classes of row 4-7 in [15, 18, Figure 1], where the rows 5-7 are not of Cartan type. They build up the tree type T_3 of [17].

Theorem 5.13 (Liftings of $\mathfrak{B}(V)$ with Cartan matrix B_2)**.** *For any lifting* A *of* $\mathfrak{B}(V)$ *as in Proposition 5.11, we have*

$$A \cong (T(V)\#\Bbbk[\Gamma])/I,$$

where I *is specified as follows:*

(1) 

(Cartan type B_2*). Let* $q_{12}q_{21} = q_{11}^{-2} = q_{22}^{-1}$.

(a) *If* $\mathrm{ord} q_{11} = 4$ *and* $q_{12} \neq \pm 1$, *then* I *is generated by*

$$[x_1x_1x_1x_2], \qquad x_1^4 - \mu_1(1 - g_1^4),$$
$$[x_1x_1x_2]^2,$$
$$[x_1x_2]^4 - \mu_{12}(1 - g_{12}^4),$$
$$x_2^2.$$

(b) *If* $\mathrm{ord} q_{11} = 4$ *and* $q_{12} = \pm 1$, *then* I *is generated by*

$$\begin{aligned}
&[x_1x_1x_1x_2], \qquad && x_1^4 - \mu_1(1-g_1^4),\\
& && [x_1x_1x_2]^2 - 8q_{11}\mu_1x_2^2 - \mu_{112}(1-g_{112}^2),\\
& && [x_1x_2]^4 - 16\mu_1x_2^4 + 4\mu_{112}q_{11}x_2^2 - \mu_{12}(1-g_{12}^4),\\
& && x_2^2 - \mu_2(1-g_2^2).
\end{aligned}$$

In both (a) *and* (b) $\dim_{\Bbbk} A = 2^2 4^2 \cdot |\Gamma| = 128 \cdot |\Gamma|$ *and a basis is*

$$\big\{x_2^{r_2}[x_1x_2]^{r_{12}}[x_1x_1x_2]^{r_{112}}x_1^{r_1}g \mid 0 \le r_1, r_{12} < 4,\ 0 \le r_2, r_{112} < 2,\ g \in \Gamma\big\}.$$

(2) 

, 

. *Let* $q_{12}q_{21} = q_{11}^{-2}$, $q_{22} = -1$.

(a) *If* $\mathrm{ord} q_{11} = 3$ *and* $q_{12} \neq \pm 1$, *then* I *is generated by*

$$\begin{aligned}
&[x_1x_1x_2x_1x_2], \qquad && x_1^3 - \mu_1(1-g_1^3),\\
& && [x_1x_2]^6 - \mu_{12}(1-g_{12}^6),\\
& && x_2^2.
\end{aligned}$$

(b) *If* $\mathrm{ord} q_{11} = 3$ *and* $q_{12} = -1$, *then* I *is generated by*

$$\begin{aligned}
&[x_1x_1x_2x_1x_2], \qquad && x_1^3,\\
& && [x_1x_2]^6 - \mu_{12}(1-g_{12}^6),\\
& && x_2^2 - \mu_2(1-g_2^2).
\end{aligned}$$

(c) *If* $\mathrm{ord} q_{11} = 3$ *and* $q_{12} = 1$, *then* I *is generated by*

$$\begin{aligned}
&[x_1x_1x_2x_1x_2] + 3\mu_1(1-q_{11})x_2^2 - \lambda_{11212}(1-g_{11212}),\\
&x_1^3 - \mu_1(1-g_1^3),\\
&[x_1x_2]^6 - s_{12} - \mu_{12}(1-g_{12}^6),\\
&x_2^2 - \mu_2(1-g_2^2),
\end{aligned}$$

where

$$\begin{aligned}
s_{12} := -3\mu_2\Big\{&(\lambda_{11212}(1-q_{11}) + 9\mu_1\mu_2q_{11})[x_1x_2]^2x_1g_2^2\\
&- q_{11}(\lambda_{11212}(1-q_{11}) + 9\mu_1\mu_2q_{11})[x_1x_2][x_1x_1x_2]g_2^2\\
&+ (\lambda_{11212}^2q_{11}^2 + 3\mu_1\mu_2\lambda_{11212}(1-q_{11}^2) - 9\mu_1^2\mu_2^2)g_1^6g_2^6\\
&+ 3\mu_1\mu_2(\lambda_{11212}(1-q_{11}^2) - 3\mu_1\mu_2)g_1^3g_2^6\\
&+ \lambda_{11212}(3\mu_1\mu_2(q_{11}-1) + \lambda_{11212})g_1^3g_2^4\\
&- 9\mu_1^2\mu_2^2g_2^6\\
&+ 3\mu_1\mu_2(\lambda_{11212}(q_{11}-1) - 9\mu_1\mu_2q_{11})g_2^4\\
&+ q_{11}(\lambda_{11212}^2 - 6\mu_1\mu_2\lambda_{11212}(1-q_{11}) - 27\mu_1^2\mu_2^2q_{11})g_2^2\Big\}.
\end{aligned}$$

In (a),(b),(c) $\dim_{\Bbbk} A = 72 \cdot |\Gamma|$. *and a basis is*

$$\big\{x_2^{r_2}[x_1x_2]^{r_{12}}[x_1x_1x_2]^{r_{112}}x_1^{r_1}g \mid 0 \le r_1 < 3,\ 0 \le r_{12} < 6,\ 0 \le r_2, r_{112} < 2,\ g \in \Gamma\big\}.$$

(d) *Let* $N := \mathrm{ord} q_{11} > 4$ *(*$N = 4$ *is* (1)*), and* $q_{12} \neq \pm 1$*. Denote*

$$N' := \mathrm{ord}(-q_{11}^{-1}) = \begin{cases} 2N, & \textit{if } N \textit{ odd}, \\ N/2, & \textit{if } N \textit{ even and } N/2 \textit{ odd}, \\ N, & \textit{if } N, N/2 \textit{ even}. \end{cases}$$

Then I *is generated by*

$$\begin{aligned} [x_1x_1x_1x_2], \qquad\qquad & x_1^N - \mu_1(1 - g_1^N), \\ & [x_1x_2]^{N'} - \mu_{12}(1 - g_{12}^{N'}), \\ & x_2^2. \end{aligned}$$

It is $\dim_{\Bbbk} A = 4NN' \cdot |\Gamma|$ *and a basis is*

$$\left\{x_2^{r_2}[x_1x_2]^{r_{12}}[x_1x_1x_2]^{r_{112}}x_1^{r_1}g \mid 0 \leq r_1 < N,\ 0 \leq r_{12} < N',\ 0 \leq r_2, r_{112} < 2,\ g \in \Gamma\right\}.$$

(3) 

, 

. *Let* $\mathrm{ord} q_{11} = 3$, $q_{12}q_{21} = q_{22}^{-1}$.

(a) *If* $q_{22} = -1$ *and* $q_{12} \neq \pm 1$*, then* I *is generated by*

$$\begin{aligned} [x_1x_1x_2x_1x_2], \qquad\qquad & x_1^3 - \mu_1(1 - g_1^3), \\ & [x_1x_1x_2]^6 - \mu_{112}(1 - g_{112}^6) \\ & x_2^2. \end{aligned}$$

(b) *If* $q_{22} = -1$ *and* $q_{12} = 1$*, then* I *is generated by*

$$\begin{aligned} [x_1x_1x_2x_1x_2], \qquad\qquad & x_1^3, \\ & [x_1x_1x_2]^6 - \mu_{112}(1 - g_{112}^6) \\ & x_2^2 - \mu_2(1 - g_2^2). \end{aligned}$$

(c) *If* $q_{22} = -1$ *and* $q_{12} = -1$*, then* I *is generated by*

$$\begin{gathered} [x_1x_1x_2x_1x_2] + 4\mu_2x_1^3g_2^2 - \lambda_{11212}(1 - g_1^3g_2^2), \\ x_1^3 - \mu_1(1 - g_1^3), \\ [x_1x_1x_2]^6 - s_{112} - \mu_{112}(1 - g_{112}^6) \\ x_2^2 - \mu_2(1 - g_2^2), \end{gathered}$$

where

$$\begin{aligned} s_{112} := -2\mu_1\Big\{ & 2(-\lambda_{11212} + 4\mu_1\mu_2)q_{11}(1 - q_{11})x_2[x_1x_1x_2]^3g_1^3g_2^2 \\ & + 2(\lambda_{11212} - 4\mu_1\mu_2)q_{11}(1 - q_{11})[x_1x_2]^2[x_1x_1x_2]^2g_1^3g_2^2 \\ & + 2(\lambda_{11212}^2 - 8\mu_1\mu_2\lambda_{11212} + 16\mu_1^2\mu_2^2)q_{11}(1 - q_{11})[x_1x_2][x_1x_1x_2]g_1^6g_2^4 \\ & + 8\mu_1\mu_2(\lambda_{11212} - 4\mu_1\mu_2)q_{11}(1 - q_{11})[x_1x_2][x_1x_1x_2]g_1^3g_2^4 \\ & + 2\lambda_{11212}(-\lambda_{11212} + 4\mu_1\mu_2)q_{11}(1 - q_{11})[x_1x_2][x_1x_1x_2]g_1^3g_2^2 \\ & + 2(-\lambda_{11212}^3 + 6\mu_1\mu_2\lambda_{11212}^2 - 16\mu_1^2\mu_2^2\lambda_{11212} + 16\mu_1^3\mu_2^3)g_1^{12}g_2^6 \\ & + (-\lambda_{11212}^3 + 12\mu_1\mu_2\lambda_{11212}^2 - 48\mu_1^2\mu_2^2\lambda_{11212} + 64\mu_1^3\mu_2^3)q_{11}(1 - q_{11})g_1^9g_2^6 \\ & + 10\mu_1\mu_2(-\lambda_{11212}^2 + 8\mu_1\mu_2\lambda_{11212} - 16\mu_1^2\mu_2^2)g_1^6g_2^6 \\ & + 2(\lambda_{11212}^3 - 7\mu_1\mu_2\lambda_{11212}^2 + 8\mu_1^2\mu_2^2\lambda_{11212} + 16\mu_1^3\mu_2^3)g_1^6g_2^4 \\ & + 16\mu_1^2\mu_2^2(\lambda_{11212} - 4\mu_1\mu_2)q_{11}(1 - q_{11})g_1^3g_2^6 \\ & + 8\mu_1\mu_2\lambda_{11212}(-\lambda_{11212} + 4\mu_1\mu_2)q_{11}(1 - q_{11})g_1^3g_2^4 \\ & + 32\mu_1^3\mu_2^3g_2^6 \end{aligned}$$

$$
\begin{aligned}
&+ \lambda_{11212}^2(\lambda_{11212} - 4\mu_1\mu_2)q_{11}(1-q_{11})g_1^3g_2^2\\
&+ 32\mu_1^2\mu_2^2(-\lambda_{11212} + \mu_1\mu_2)g_2^4\\
&+ 4\mu_1\mu_2(3\lambda_{11212}^2 - 8\mu_1\mu_2\lambda_{11212} + 8\mu_1^2\mu_2^2)g_2^2\Big\}
\end{aligned}
$$

In (a),(b),(c) $\dim_{\Bbbk} A = 108 \cdot |\Gamma|$ *and a basis is*

$$\big\{x_2^{r_2}[x_1x_2]^{r_{12}}[x_1x_1x_2]^{r_{112}}x_1^{r_1}g \mid 0 \le r_1, r_{12} < 3,\ 0 \le r_2 < 2,\ 0 \le r_{112} < 6,\ g \in \Gamma\big\}.$$

(4) 

, 

. *Let* $\mathrm{ord} q_{11} = 3$, $q_{12}q_{21} = -q_{11}$ *of order* 6, $q_{22} = -1$.

(a) *If* $q_{12} \neq \pm 1$, *then* I *is generated by*

$$
\begin{aligned}
&[x_1x_1x_2x_1x_2], && x_1^3 - \mu_1(1-g_1^3),\\
& && x_2^2.
\end{aligned}
$$

(b) *If* $q_{12} = 1$, *then* I *is generated by*

$$
\begin{aligned}
&[x_1x_1x_2x_1x_2], && x_1^3,\\
& && x_2^2 - \mu_2(1-g_2^2).
\end{aligned}
$$

(c) *If* $q_{12} = -1$, *then* I *is generated by*

$$
\begin{gathered}
[x_1x_1x_2x_1x_2] - \mu_2(1+q_{11})x_1^3g_2^2 - \lambda_{11212}(1-g_{11212}),\\
x_1^3 - \mu_1(1-g_1^3),\\
x_2^2 - \mu_2(1-g_2^2).
\end{gathered}
$$

In (a),(b),(c) $\dim_{\Bbbk} A = 36 \cdot |\Gamma|$ *and a basis is*

$$\big\{x_2^{r_2}[x_1x_2]^{r_{12}}[x_1x_1x_2]^{r_{112}}x_1^{r_1}g \mid 0 \le r_1, r_{12} < 3,\ 0 \le r_2, r_{112} < 2,\ g \in \Gamma\big\}.$$

Proof. We proceed as in the proof of Theorem 5.9.

- $(T(V)\#\Bbbk[\Gamma])/I$ is a Hopf algebra, since I is generated by skew-primitive elements: Again the elements $x_i^{N_i} - \mu_i(1-g_i^{N_i})$ and $[x_1x_1x_1x_2] - \lambda_{1112}(1-g_{1112})$ are skew-primitive if $q_{12}q_{21} = q_{11}^{-2}$ by Lemma 4.2.

(1a) By Lemma 4.3(1) $[x_1x_2]^4 \in P_{g_{12}^4}^{\chi_{12}^4}$ and hence also $[x_1x_2]^4 - \mu_{12}(1-g_{12}^4) \in P_{g_{12}^4}^{\chi_{12}^4}$. A direct computation yields $[x_1x_1x_2]^2 \in P_{g_{112}^2}^{\chi_{112}^2}$.

(1b) Again direct computation shows that $[x_1x_1x_2]^2 - 8q_{11}\mu_1x_2^2 - \mu_{112}(1-g_{112}^2)$ and $[x_1x_2]^4 - 16\mu_1x_2^4 + 4\mu_{112}q_{11}x_2^2 - \mu_{12}(1-g_{12}^4)$ are skew primitive; we used the computer algebra system FELIX [9].

(2a) We have $[x_1x_1x_2x_1x_2] \in P_{g_{11212}}^{\chi_{11212}}$ by Lemma 4.4(2). Further $[x_1x_2]^6 \in P_{g_{12}^6}^{\chi_{12}^6}$ by Lemma 4.3(1).

(2b) Again $[x_1x_1x_2x_1x_2] \in P_{g_{11212}}^{\chi_{11212}}$ by Lemma 4.4(2) and a direct computation yields $[x_1x_2]^6 - \mu_{12}(1-g_{12}^6) \in P_{g_{12}^6}^{\chi_{12}^6}$.

(2c) Using FELIX we get that all elements are skew-primitive.

(2d) This is again Lemma 4.3(1).

(3a) and (3b): $[x_1x_1x_2x_1x_2] \in P_{g_{11212}}^{\chi_{11212}}$ by Lemma 4.4(1). Straightforward calculation shows that $[x_1x_1x_2]^6 - \mu_{112}(1-g_{112}^6) \in P_{g_{112}^6}^{\chi_{112}^6}$; here again we used FELIX.

(3c) is computed using FELIX.

(4a) and (4b): $[x_1x_1x_2x_1x_2] \in P_{g_{11212}}^{\chi_{11212}}$ by Lemma 4.4(1).

(4c) Looking at the coproduct computed in Lemma 4.4(1) we deduce that the element $[x_1x_1x_2x_1x_2] - \mu_2(1+q_{11})x_1^3g_2^2$ and hence $[x_1x_1x_2x_1x_2] - \mu_2(1+q_{11})x_1^3g_2^2 - \lambda_{11212}(1-g_{11212})$ is skew-primitive.

- The statement on the basis and dimension of $(T(V)\#\Bbbk[\Gamma])/I$ is proven in [23].
- $(T(V)\#\Bbbk[\Gamma])/I$ is pointed by the same argument as in the proof of Theorem 5.9.
- The surjective Hopf algebra map as given in the proof of Theorem 5.9

$$T(V)\#\Bbbk[\Gamma] \to \operatorname{gr}((T(V)\#\Bbbk[\Gamma])/I)$$

factorizes to an isomorphism $\mathfrak{B}(V)\#\Bbbk[\Gamma] \xrightarrow{\sim} \operatorname{gr}((T(V)\#\Bbbk[\Gamma])/I)$: Again we look at the coradical filtration. All equations of I are of the form $[w]-c_w$, $[u]^{N_u}-d_u$ with c_w resp. d_u of lower degree in $\operatorname{gr}((T(V)\#\Bbbk[\Gamma])/I)$, hence $[w]=0$, $[u]^{N_u}=0$ in $\operatorname{gr}((T(V)\#\Bbbk[\Gamma])/I)$.

- Like before, for a lifting A we have to check whether the surjective Hopf algebra map

$$T(V)\#\Bbbk[\Gamma] \to A$$

which takes x_i to a_i and g to g factorizes to

$$(T(V)\#\Bbbk[\Gamma])/I \xrightarrow{\sim} A.$$

By Lemma 5.6 the relations concerning the elements $a_i^{N_i}$ and $[a_1a_1a_1a_2]$ are of the right form. We deduce from Lemma 5.2 that the relations also hold in A: this is just combinatorics on the braiding matrices which we want to demonstrate for the following.

(1a) We have $\chi_{112}^2 \neq \varepsilon$ by Lemma 5.2(2a), since $q_{1,112}^2 = q_{12}^2 \neq 1$. Further $P_{g_{112}^2}^{\chi_{112}^2} = 0$ by Lemma 5.2(2b): Suppose $q_{21}^4q_{22}^2 = q_{21}$, $q_{12}^4q_{22}^2 = q_{12}$, then $q_{21}^3 = q_{12}^3 = 1$, which contradicts $q_{12}q_{21} = q_{11}^{-2} = -1$; also if $q_{11}^4q_{12}^2 = q_{12}$, $q_{11}^4q_{21}^2 = q_{21}$, then $q_{12}=q_{21}=1$, again a contradiction to $q_{12}q_{21} = q_{11}^{-2} = -1$. Hence $[a_1a_1a_2]^2 = 0$. The other cases work in exactly the same manner. □

Remark 5.14. The Conjecture 5.4 is true in the above cases. Further note that in (2c) $s_{12} \notin \Bbbk[\Gamma]$ and in (3c) $s_{112} \notin \Bbbk[\Gamma]$.

Further we want to note the cases not treated in the theorem above:

1. The case (1) when $5 \neq N := \operatorname{ord} q_{11} \geq 3$ is odd is treated in [10], and the case $N=5$ in [11].
2. There is no general method for (1) in the case $N := \operatorname{ord} q_{11} \geq 6$ is even. Here $\operatorname{ord} q_{22} = \operatorname{ord} q_{11}^2 = \frac{N}{2}$.
3. There is no general method for (2d) in the case $q_{12} = \pm 1$.
4. There is no general method for (3) in the case $N := \operatorname{ord} q_{22} \geq 4$. The case $N=3$ is (1) of the theorem above or (1) of Theorem 5.9.

5.5 Lifting of $\mathfrak{B}(V)$ of non-standard type

In this section we want to lift some of the Nichols algebras of the Weyl equivalence classes of rows 8 and 9 of [15, 18, Figure 1] which are not of standard type, namely for $\operatorname{ord}\zeta = 12$ we lift

$$\overset{-\zeta^{-2}}{\circ}\overset{-\zeta^{3}}{\text{———}}\overset{-\zeta^{2}}{\circ},\quad \overset{-\zeta^{-2}}{\circ}\overset{\zeta^{-1}}{\text{———}}\overset{-1}{\circ},\quad \overset{-\zeta^{2}}{\circ}\overset{-\zeta}{\text{———}}\overset{-1}{\circ},\quad \overset{-\zeta^{3}}{\circ}\overset{\zeta}{\text{———}}\overset{-1}{\circ},\quad \overset{-\zeta^{3}}{\circ}\overset{-\zeta^{-1}}{\text{———}}\overset{-1}{\circ}$$

of row 8, and

$$\overset{-\zeta^{2}}{\circ}\overset{\zeta^{3}}{\text{———}}\overset{-1}{\circ},\quad \overset{-\zeta^{-1}}{\circ}\overset{-\zeta^{3}}{\text{———}}\overset{-1}{\circ}$$

of row 9. Again, at first we give a nice presentation of the ideal cancelling the redundant relations of the ideals given in [17]:

Proposition 5.15 (Nichols algebras of rows 8 and 9)**.** *The following finite-dimensional Nichols algebras $\mathfrak{B}(V)$ with braiding matrix (q_{ij}) of rows 8 and 9 of [15, 18, Figure 1] are represented as follows: Let $\zeta \in \Bbbk^{\times}$, $\operatorname{ord}\zeta = 12$.*

(1) $\overset{-\zeta^{-2}}{\circ}\overset{-\zeta^{3}}{\text{———}}\overset{-\zeta^{2}}{\circ}$. *Let $q_{11} = -\zeta^{-2}$, $q_{12}q_{21} = -\zeta^{3}$, $q_{22} = -\zeta^{2}$, then*

$$\mathfrak{B}(V) = T(V)/\big([x_1x_1x_2x_2] - \frac{1}{2}q_{11}q_{12}(q_{12}q_{21} - q_{11})(1 - q_{12}q_{21})[x_1x_2]^2, x_1^3,\ x_2^3\big)$$

with basis $\big\{x_2^{r_2}[x_1x_2x_2]^{r_{122}}[x_1x_2]^{r_{12}}[x_1x_1x_2]^{r_{112}}x_1^{r_1} \mid 0 \le r_1, r_2 < 3,\ 0 \le r_{112}, r_{122} < 2,\ 0 \le r_{12} < 4\big\}$ *and* $\dim_{\Bbbk}\mathfrak{B}(V) = 144$.

(2) $\overset{-\zeta^{-2}}{\circ}\overset{\zeta^{-1}}{\text{———}}\overset{-1}{\circ}$, $\overset{-\zeta^{2}}{\circ}\overset{-\zeta}{\text{———}}\overset{-1}{\circ}$. *Let $q_{11} = -\zeta^{2}$, $q_{12}q_{21} = -\zeta$, $q_{22} = -1$, or $q_{11} = -\zeta^{-2}$, $q_{12}q_{21} = \zeta^{-1}$, $q_{22} = -1$, then*

$$\mathfrak{B}(V) = T(V)/\big([x_1x_1x_2x_1x_2x_1x_2],\ x_1^3,\ x_2^2\big)$$

with basis $\big\{x_2^{r_2}[x_1x_2]^{r_{12}}[x_1x_1x_2x_1x_2]^{r_{11212}}[x_1x_1x_2]^{r_{112}}x_1^{r_1} \mid 0 \le r_1, r_{112} < 3,\ 0 \le r_2, r_{11212} < 2,\ 0 \le r_{12} < 4\big\}$ *and* $\dim_{\Bbbk}\mathfrak{B}(V) = 144$.

(3) $\overset{-\zeta^{3}}{\circ}\overset{\zeta}{\text{———}}\overset{-1}{\circ}$, $\overset{-\zeta^{3}}{\circ}\overset{-\zeta^{-1}}{\text{———}}\overset{-1}{\circ}$. *Let $q_{11} = -\zeta^{3}$, $q_{12}q_{21} = \zeta$, $q_{22} = -1$, or $q_{11} = -\zeta^{3}$, $q_{12}q_{21} = -\zeta^{-1}$, $q_{22} = -1$, then*

$$\mathfrak{B}(V) = T(V)/\big([x_1x_1x_2x_1x_2],\ x_1^4,\ x_2^2\big)$$

with basis $\big\{x_2^{r_2}[x_1x_2]^{r_{12}}[x_1x_1x_2]^{r_{112}}[x_1x_1x_1x_2]^{r_{1112}}x_1^{r_1} \mid 0 \le r_1 < 4,\ 0 \le r_{12}, r_{112} < 3,\ 0 \le r_2, r_{1112} < 2\big\}$ *and* $\dim_{\Bbbk}\mathfrak{B}(V) = 144$.

(4) $\overset{-\zeta^{2}}{\circ}\overset{\zeta^{3}}{\text{———}}\overset{-1}{\circ}$. *Let $q_{11} = -\zeta^{2}$, $q_{12}q_{21} = \zeta^{3}$, $q_{22} = -1$, then*

$$\mathfrak{B}(V) = T(V)/\big([x_1x_1x_2x_1x_2x_1x_2],\ x_1^3,\ [x_1x_2]^{12},\ x_2^2\big)$$

with basis $\big\{x_2^{r_2}[x_1x_2]^{r_{12}}[x_1x_1x_2x_1x_2]^{r_{11212}}[x_1x_1x_2]^{r_{112}}x_1^{r_1} \mid 0 \le r_1, r_{112} < 3,\ 0 \le r_2, r_{11212} < 2,\ 0 \le r_{12} < 12\big\}$ *and* $\dim_{\Bbbk}\mathfrak{B}(V) = 432$.

(5) $\overset{-\zeta^{-1}}{\circ}\overset{-\zeta^{3}}{\text{———}}\overset{-1}{\circ}$. *Let $q_{11} = -\zeta^{-1}$, $q_{12}q_{21} = -\zeta^{3}$, $q_{22} = -1$, then*

$$\mathfrak{B}(V) = T(V)/\big([x_1x_1x_1x_1x_2],\ [x_1x_1x_2x_1x_2],\ x_1^{12},\ x_2^2\big)$$

with basis $\big\{x_2^{r_2}[x_1x_2]^{r_{12}}[x_1x_1x_2]^{r_{112}}[x_1x_1x_1x_2]^{r_{1112}}x_1^{r_1} \mid 0 \le r_1 < 12,\ 0 \le r_{12}, r_{112} < 3,\ 0 \le r_2, r_{1112} < 2\big\}$ *and* $\dim_{\Bbbk}\mathfrak{B}(V) = 432$.

Remark 5.16. The Nichols algebras of Proposition 5.15 have different PBW bases, also if they are in the same Weyl equivalence class. They build up the tree types T_4, T_5 and T_7 of [17].

Theorem 5.17 (Liftings of $\mathfrak{B}(V)$ of rows 8 and 9)**.** *For any lifting A of $\mathfrak{B}(V)$ as in Proposition 5.15, we have*

$$A \cong (T(V)\#\Bbbk[\Gamma])/I,$$

where I is specified as follows: Let $\zeta \in \Bbbk^\times$, $\mathrm{ord}\zeta = 12$.

(1) 

. *Let $q_{11} = -\zeta^{-2}$, $q_{12}q_{21} = -\zeta^3$, $q_{22} = -\zeta^2$.*

(a) *If $q_{12}^3 \neq 1$, then I is generated by*

$$\begin{aligned}&[x_1x_1x_2x_2] - \frac{1}{2}q_{11}q_{12}(q_{12}q_{21} - q_{11})(1 - q_{12}q_{21})[x_1x_2]^2,\\ &x_1^3 - \mu_1(1 - g_1^3),\\ &x_2^3.\end{aligned}$$

(b) *If $q_{12}^3 = 1$, then I is generated by*

$$\begin{aligned}&[x_1x_1x_2x_2] - \frac{1}{2}q_{11}q_{12}(q_{12}q_{21} - q_{11})(1 - q_{12}q_{21})[x_1x_2]^2,\\ &x_1^3,\\ &x_2^3 - \mu_2(1 - g_2^3).\end{aligned}$$

In (a),(b) $\dim_\Bbbk A = 144 \cdot |\Gamma|$ *and a basis is*

$$\big\{x_2^{r_2}[x_1x_2x_2]^{r_{122}}[x_1x_2]^{r_{12}}[x_1x_1x_2]^{r_{112}}x_1^{r_1}g \mid 0 \le r_1, r_2 < 3,\\ 0 \le r_{112}, r_{122} < 2,\ 0 \le r_{12} < 4,\ g \in \Gamma\big\}.$$

(2) 

, 

. *Let $q_{11} = -\zeta^2$, $q_{12}q_{21} = -\zeta$, $q_{22} = -1$, or $q_{11} = -\zeta^{-2}$, $q_{12}q_{21} = \zeta^{-1}$, $q_{22} = -1$.*

(a) *If $q_{12} \neq \pm 1$, then I is generated by*

$$[x_1x_1x_2x_1x_2x_1x_2], \qquad x_1^3 - \mu_1(1 - g_1^3),\quad x_2^2.$$

(b) *If $q_{12} = \pm 1$, then I is generated by*

$$\begin{aligned}&[x_1x_1x_2x_1x_2x_1x_2] + \mu_2 q_{12}(q_{11}q_{12}q_{21} + q_{12}q_{21} - 1)[x_1x_1x_2]x_1^2g_2^2,\\ &x_1^3,\\ &x_2^2 - \mu_2(1 - g_2^2).\end{aligned}$$

In (a),(b) $\dim_\Bbbk A = 144 \cdot |\Gamma|$ *and a basis is*

$$\big\{x_2^{r_2}[x_1x_2]^{r_{12}}[x_1x_1x_2x_1x_2]^{r_{11212}}[x_1x_1x_2]^{r_{112}}x_1^{r_1}g \mid 0 \le r_1, r_{112} < 3,\\ 0 \le r_2, r_{11212} < 2, 0 \le r_{12} < 4,\ g \in \Gamma\big\}.$$

(3) $\overset{-\zeta^{3}}{\circ}\overset{\zeta}{\rule{2em}{0.4pt}}\overset{-1}{\circ}$, $\overset{-\zeta^{3}}{\circ}\overset{-\zeta^{-1}}{\rule{2em}{0.4pt}}\overset{-1}{\circ}$. *Let* $q_{11} = -\zeta^3$, $q_{12}q_{21} = \zeta$, $q_{22} = -1$, *or* $q_{11} = -\zeta^3$, $q_{12}q_{21} = -\zeta^{-1}$, $q_{22} = -1$.

(a) *If* $q_{12} \neq \pm 1$, *then* I *is generated by*

$$\begin{aligned}&[x_1x_1x_2x_1x_2], && x_1^4 - \mu_1(1-g_1^3),\\ & && x_2^2.\end{aligned}$$

(b) *If* $q_{12} = \pm 1$, *then* I *is generated by*

$$\begin{aligned}&[x_1x_1x_2x_1x_2] - \mu_2 q_{12}(q_{11} + 2q_{12}^2q_{21}^2 - q_{12}q_{21})x_1^3g_2^2,\\ &x_1^4,\\ &x_2^2 - \mu_2(1-g_2^2).\end{aligned}$$

In (a),(b) $\dim_{\Bbbk} A = 144 \cdot |\Gamma|$ *and a basis is*

$$\big\{x_2^{r_2}[x_1x_2]^{r_{12}}[x_1x_1x_2]^{r_{112}}[x_1x_1x_1x_2]^{r_{1112}}x_1^{r_1}g \mid 0 \le r_1 < 4,\\ 0 \le r_{12}, r_{112} < 3,\ 0 \le r_2, r_{1112} < 2,\ g \in \Gamma\big\}.$$

(4) $\overset{-\zeta^{2}}{\circ}\overset{\zeta^{3}}{\rule{2em}{0.4pt}}\overset{-1}{\circ}$. *Let* $q_{11} = -\zeta^2$, $q_{12}q_{21} = \zeta^3$, $q_{22} = -1$.

(a) *If* $q_{12} \neq \pm 1$, *then* I *is generated by*

$$\begin{aligned}&[x_1x_1x_2x_1x_2x_1x_2], && x_1^3 - \mu_1(1-g_1^3),\\ & && [x_1x_2]^{12} - \mu_{12}(1-g_{12}^{12}),\\ & && x_2^2.\end{aligned}$$

It is $\dim_{\Bbbk} A = 432 \cdot |\Gamma|$ *and a basis is*

$$\big\{x_2^{r_2}[x_1x_2]^{r_{12}}[x_1x_1x_2x_1x_2]^{r_{11212}}[x_1x_1x_2]^{r_{112}}x_1^{r_1}g \mid 0 \le r_1, r_{112} < 3,\\ 0 \le r_2, r_{11212} < 2,\ 0 \le r_{12} < 12,\ g \in \Gamma\big\}$$

(b) (incomplete) $q_{12} = \pm 1$, *then* I *is generated by*

$$\begin{aligned}&[x_1x_1x_2x_1x_2x_1x_2] + q_{12}2\mu_2(q_{12}q_{21}+1)[x_1x_1x_2]x_1^2g_2^2,\\ &x_1^3,\\ &[x_1x_2]^{12} - d_{12},\\ &x_2^2 - \mu_2(1-g_1^2).\end{aligned}$$

(5) $\overset{-\zeta^{-1}}{\circ}\overset{-\zeta^{3}}{\rule{2em}{0.4pt}}\overset{-1}{\circ}$. *Let* $q_{11} = -\zeta^{-1}$, $q_{12}q_{21} = -\zeta^3$, $q_{22} = -1$.

(a) *If* $q_{12} \neq \pm 1$, *then* I *is generated by*

$$\begin{aligned}&[x_1x_1x_1x_1x_2], && x_1^{12} - \mu_1(1-g_1^{12}),\\ &[x_1x_1x_2x_1x_2], && x_2^2.\end{aligned}$$

(b) *If* $q_{12} = \pm 1$, *then* I *is generated by:*

$$\begin{aligned}&[x_1x_1x_1x_1x_2], && x_1^{12} - \mu_1(1-g_1^{12}),\\ &[x_1x_1x_2x_1x_2] + 2\mu_2q_{12}x_1^3g_2^2, && x_2^2 - \mu_2(1-g_2^2).\end{aligned}$$

In (a),(b) $\dim_{\Bbbk} A = 432 \cdot |\Gamma|$ *and a basis is*

$$\big\{x_2^{r_2}[x_1x_2]^{r_{12}}[x_1x_1x_2]^{r_{112}}[x_1x_1x_1x_2]^{r_{1112}}x_1^{r_1}g \mid 0 \le r_1 < 12,\\ 0 \le r_{12}, r_{112} < 3,\ 0 \le r_2, r_{1112} < 2,\ g \in \Gamma\big\}.$$

Proof. We argue exactly as in the proofs of Theorem 5.9 and 5.13.

- $(T(V)\#\Bbbk[\Gamma])/I$ is a Hopf algebra, since I is generated by skew-primitive elements: The elements $x_i^{N_i} - \mu_i(1-g_i^{N_i})$ and $[x_1x_1x_1x_1x_2] - \lambda_{11112}(1-g_{11112})$ are skew-primitive if $q_{12}q_{21} = q_{11}^{-2}$ by Lemma 4.2. For the elements $[x_1x_1x_2x_1x_2] - c_{11212}$ and $[x_1x_1x_2x_1x_2x_1x_2] - c_{1121212}$ we use Lemma 4.4 and for $[x_1x_2]^{N_{12}} - d_{12}$ Lemma 4.3(1). Further in (1) $[x_1x_1x_2x_2] - \frac{1}{2}q_{11}q_{12}(q_{12}q_{21} - q_{11})(1 - q_{12}q_{21})[x_1x_2]^2$ is skew-primitive by a straightforward calculation.
- The statement on the basis and dimension of $(T(V)\#\Bbbk[\Gamma])/I$ is proven in [23].
- $(T(V)\#\Bbbk[\Gamma])/I$ is pointed and $\operatorname{gr}((T(V)\#\Bbbk[\Gamma])/I) \cong \mathfrak{B}(V)\#\Bbbk[\Gamma]$ by the same arguments as in the proofs of Theorems 5.9 and 5.13.
- Also in the same way, the surjective Hopf algebra map $T(V)\#\Bbbk[\Gamma] \to A$ factorizes to an isomorphism

$$(T(V)\#\Bbbk[\Gamma])/I \xrightarrow{\sim} A$$

by Lemma 5.6 and 5.2, doing the combinatorics on the braiding matrices. □

Remark 5.18. The Conjecture 5.4 is true in the above cases. Further note that in (1) $r_{1122} \notin \Bbbk[\Gamma]$ (as well as $r_{1122} \neq 0$ in $\mathfrak{B}(V)$), in (2b) $r_{1121212} \notin \Bbbk[\Gamma]$, in (3b) $r_{11212} \notin \Bbbk[\Gamma]$, in (4b) $r_{1121212} \notin \Bbbk[\Gamma]$ and in (5b) $r_{11212} \notin \Bbbk[\Gamma]$.

References

[1] N. Andruskiewitsch and I. Angiono. On Nichols algebras with generic braiding. In I. Shestakov P.F. Smith T. Brzezinski, J.L. Gómez Pardo, editor, *Modules and Comodules*, Trends in Mathematics, pages 47–64, 2008.

[2] N. Andruskiewitsch, I. Heckenberger, and H.-J. Schneider. The Nichols algebra of a semisimple Yetter-Drinfeld module, 2008. preprint `arXiv.org:0803.2430`.

[3] N. Andruskiewitsch and H.-J. Schneider. Lifting of quantum linear spaces and pointed Hopf algebras of order p^3. *J. Algebra*, 209:658–691, 1998.

[4] N. Andruskiewitsch and H.-J. Schneider. Finite quantum groups and Cartan matrices. *Adv. in Math.*, 154:1–45, 2000.

[5] N. Andruskiewitsch and H.-J. Schneider. Lifting of Nichols algebras of type A_2 and pointed Hopf algebras of order p^4. In S. Caenepeel and F. van Oystaeyen, editors, *Hopf algebras and quantum groups: Proceedings of the Brussels Conference*, volume 209 of *Lecture Notes in Pure and Appl. Math.*, pages 1–14. Marcel Dekker, 2000.

[6] N. Andruskiewitsch and H.-J. Schneider. Pointed Hopf algebras. In *New directions in Hopf algebras*, volume 43, pages 1–68. MSRI Publications, Cambridge Univ. Press, 2002.

[7] N. Andruskiewitsch and H.-J. Schneider. On the classification of finite-dimensional pointed Hopf algebras. 2007. to appear in *Ann. Math.*, `arXiv math.QA/0502157`.

[8] I. Angiono. On Nichols algebras with standard braiding. 2008. preprint `arXive math.QA/0804.0816`.

[9] J. Apel and U. Klaus. FELIX, a special computer algebra system for the computation in commutative and non-commutative rings and modules. available at `http://felix.hgb-leipzig.de/`.

[10] M. Beattie, S. Dăscălescu, and S. Raianu. Lifting of Nichols algebras of type B_2. *Israel J. Math.*, 132:1–28, 2002.

[11] D. Didt. Linkable Dynkin diagrams and quasi-isomorphisms for finite dimensional pointed Hopf algebras. 2003.

[12] M. Graña. On pointed Hopf algebras of dimension p^5. *Glasgow Math. J.*, 42:405–419, 2000.

[13] M. Graña. Pointed Hopf algebras of dimension 32. *Communications in Algebra*, 28:2935–2976, 2000.

[14] I. Heckenberger. Classification of arithmetic root systems of rank 3. *Actas del "XVI Coloquio Latinoamericano de Álgebra"*, pages 227–252, 2005. Colonia, Uruguay.

[15] I. Heckenberger. Weyl equivalence for rank 2 Nichols algebras of diagonal type. *Ann. Univ. Ferrara - Sez. VII - Sc. Mat.*, LI:281–289, 2005.

[16] I. Heckenberger. The Weyl groupoid of a Nichols algebra of diagonal type. *Invent. Math.*, 164:175–188, 2006.

[17] I. Heckenberger. Examples of finite dimensional rank 2 Nichols algebras of diagonal type. *Compositio Math.*, 2007.

[18] I. Heckenberger. Rank 2 Nichols algebras with finite arithmetic root system. *Algebr. Represent. Theor.*, 11:115–132, 2008.

[19] I. Heckenberger. Classification of arithmetic root systems. *Adv. Math.*, 220:59–124, 2009.

[20] I. Heckenberger and H.-J. Schneider. Root systems and Weyl groupoids for Nichols algebras, 2008. preprint `arXiv.org:0807.0691`.

[21] M. Helbig. *Lifting of Nichols algebras.* Südwestdeutscher Verlag für Hochschulschriften, 2009. available at `http://edoc.ub.uni-muenchen.de/10378/`.

[22] M. Helbig. On the presentation of pointed Hopf algebras. preprint, available at `http://arxiv.org/`, 2010.

[23] M. Helbig. A PBW basis criterion for pointed Hopf algebras. preprint, available at `http://arxiv.org/`, 2010.

[24] V. Kharchenko. A quantum analog of the Poincaré-Birkhoff-Witt theorem. *Algebra and Logic*, 38:259–276, 1999.

[25] M. Lothaire. *Combinatorics on Words*, volume 17 of *Encyclopedia of Mathematics.* Addison-Wesley, 1983.

[26] S. Montgomery. *Hopf Algebras and Their Action on Rings*. CBSM Regional Conference Series in Mathematics, Vol. 82, American Mathematical Society, 1993.

[27] D. Radford. Finite-dimensional simple-pointed Hopf algebras. *J. Algebra*, 211:686–710, 1999.

[28] C. Reutenauer. *Free Lie Algebras*, volume 7 of *London Mathematical Society Monographs, New Series*. Clarendon Press, London, 1993.

[29] S. Ufer. PBW bases for a class of braided Hopf algebras. *J. Algebra*, 280:84–119, 2004.